

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 63/00, 43/04, C12N 15/00, C07H 21/02	A1	(11) International Publication Number: WO 99/26480 (43) International Publication Date: 3 June 1999 (03.06.99)
(21) International Application Number: PCT/US98/24950 (22) International Filing Date: 20 November 1998 (20.11.98) (30) Priority Data: 08/975,424 20 November 1997 (20.11.97) US (71) Applicants: GENETIX PHARMACEUTICALS, INC. [US/US]; 840 Memorial Drive, Cambridge, MA 02139 (US). MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02139 (US). (72) Inventors: LEBOULCH, Philippe; Flagship Warf, Unit 729, 197 8th Street, Charlestown, MA 02129 (US). PAWLIUK, Robert, James; Apartment 3, 52 Maple Avenue, Cambridge, MA 02129 (US). BACHELOT, Thomas; 53, rue Pierre Brunier, F-69300 Caluire (FR). (74) Agent: CLARK, Paul, T.; Clark & Elbing LLP, 176 Federal Street, Boston, MA 02110-2214 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: ANTI-ANGIOGENIC GENE THERAPY VECTORS AND THEIR USE IN TREATING ANGIOGENESIS-RELATED DISEASES (57) Abstract A method for inhibiting tumor growth in a human patient harboring a solid tumor, said method comprising administering to said patient a nucleic acid molecule which expresses in said patient an anti-angiogenic polypeptide selected from the group consisting of human angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fragments thereof, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

15

25

In a second, related aspect, the invention features tumor inhibition, of

-2-

the type just described, using nucleic acids molecules of the formula A-B, where A and B are polypeptide and/or export signal joined by a peptide bond; peptide A contains at least 100 amino acids and includes at least kringle 1, 2, and 3 of human or murine angiostatin; and peptide B contains at least 100 amino acids and includes at least 75% of the amino acid sequence of human or murine endostatin. Expression of the fusion anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth. In some embodiments of this hybrid polypeptide and/or export signal method, polypeptide and/or export signal A further includes kringle region 4 of angiostatin, and can also include kringle region 5 of plasminogen (the larger protein molecule of which angiostatin is a portion).

In both aspects of the invention, the nucleic acid molecule preferably constitutes a portion of a viral vector or a plasmid, which can either be administered to the patient so that cells of the patient in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells are infected or transfected with the nucleic acid encoding the angiogenesis-inhibiting polypeptide, or cells (of the patient, or another human donor, or an animal) are infected or transfected *ex-vivo*, and those infected or transfected cells are then infused into the patient so that the anti-angiogenic polypeptide is expressed in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells.

As will be discussed in more detail below, in particularly effective embodiments, the nucleic acid molecule includes a nucleotide sequence

-3-

encoding a preactivation polypeptide and/or export signal for effecting Golgi and/or endoplasmic reticulum export of the anti-angiogenic polypeptide.

In another aspect, the invention features a method for treating a human patient suffering from diabetic retinopathy, involving administering to
5 the patient one of the nucleic acid molecules described above.

The above and other features, objects and advantages of the present invention will be better understood by a reading of the following specification in conjunction with the drawings.

Brief Description of the Drawings

10 Fig. 1 depicts the structural relationship of angiostatin with plasminogen.

Fig. 2 depicts the structural relationship of endostatin with collagen type XVIII.

Fig. 3 depicts various viral (A. MSCV murine retrovirus; B. Adeno-associated virus; C. HIV based retrovirus; E. recombinant adeno-virus) and
15 non-viral (D. plasmid) vectors used in the construction of gene therapy vectors for this invention.

References:

- A. Hawley, R.G., F. H. L. Lieu, A. Z. C. Fong, and T. S. Hawley. 1994. Versatile retroviral vectors for potential use in gene therapy. *Gene Therapy* 1:136.
- B. Hargrove, P. H., E. F. Vanin, G. J. Kurtzman and A. W. Nienhuis. 1997. High-level globin gene expression mediated by a recombinant adeno-associated virus genome that contains a 3' γ globin gene regulatory element and integrates as tandem copies in erythroid cells. *Blood* 89:2167.
- C. Naldini, L., U. Blomer, P. Gallay, D. Ory, R. Mulligan, F. H. Gage, I. M. Verma, and D. Trono. 1996. In vivo gene delivery and stable transduction of nondividing cells by a Lentiviral vector. *Science* 272: 263.
- D.
- E. Ohashi, T., K. Watabe, K. Uehara, W. S. Sly, C. Vogler, and Y. Eto. 1997. Adenovirus-mediated gene transfer and expression of human β -glucuronidase gene in the liver, spleen, and central nervous system in mucopolysaccharide type VII mice. *PNAS* 94: 1287.

MSCV: Murine Stem Cell Virus
 LTR: Long Terminal Repeat
 RSV: Rous Sarcoma Virus
 ITR: Inverted Terminal Repeat
 HIV: Human Immunodeficiency Virus
 IRES: Internal Ribosomal Entry Site
 GFP: Green Fluorescence Protein
 HBPRE: Hepatitis B Export Element
 RRE: Rev Response Element
 polyA: polyadenylation site
 Ψ +: , , viral packaging sequence

The inverted triangle shows the site at which the anti-angiogenic constructs will be inserted using engineered MluI and XhoI restriction sites.

* denotes specific mutations within the long terminal repeat and leader which bestows the ability for expression in embryonic stem and hematopoietic stem cells.

The arrow denotes the direction of transcription.

Fig. 4 depicts in the left (A) panel nude mice which were implanted with human neuroblastoma cells (line SK-N-AS) transduced with a mock virus and in the right (B) panel, nude mice which were transplanted with human neuroblastoma cells transduced with a retroviral gene therapy vector encoding an angiostatin-endostatin fusion protein.

Fig. 5 shows the nucleotide sequence (SEQ ID NO: 1) and amino

-5-

acid sequence (SEQ ID NO: 2) of human plasminogen and the nucleotide sequence (SEQ ID NO: 5) and amino acid sequence (SEQ ID NO: 6) of human angiostatin.

Fig. 6 shows the nucleotide sequence (SEQ ID NO: 9) and amino acid sequence (SEQ ID NO: 10) of murine endostatin.

Fig. 7 shows the nucleotide sequence (SEQ ID NO: 3) and amino acid sequence (SEQ ID NO: 4) of murine plasminogen and the nucleotide (SEQ ID NO: 7) and amino acid sequence (SEQ ID NO: 8) of murine angiostatin.

Detailed Description

10 This invention provides gene therapy using a vector having a nucleotide sequence encoding one of the above-identified anti-angiogenic polypeptides. Described below in more detail are some of the components of the vectors and methods of the invention.

By a gene therapy vector is meant a vector useful for gene therapy.

15 Gene therapy vectors carry a gene of interest that is useful for gene therapy. The gene therapy vectors are able to be transferred to the cells of an animal, e.g., a human, and are able to express the gene of interest in such cells so as to effect gene therapy. The vector can be, e.g., chromosomal, non-chromosomal, or synthetic, and can be RNA or DNA. The vector can be, e.g., a plasmid, a virus or a phage. Preferred vectors include, e.g., retroviral vectors, adenoviral vectors, adeno-associated vectors, herpes virus vectors, Simliki Forest Virus-based vector, Human Immunodeficiency virus, Simian Immunodeficiency virus, and non-viral plasmids. A preferred retroviral vector is Murine Stem Cell Virus (MSCV), which is a variant of Moloney Murine Leukemia Virus

20 (MoMLV).

By anti-angiogenic polypeptide is meant a polypeptide which inhibits

-6-

angiogenesis. The terms polypeptide, protein and polypeptide and/or export signal are used interchangeably herein. By angiogenesis is meant the process by which new vasculature, in particular, new capillaries, are formed from existing vasculature. Angiogenesis is a complex process entailing numerous steps, including local dissolution of the basement membrane, migration of endothelial cells into the surrounding stroma, proliferation of the endothelial cells at the leading edge to form a migrating column of cells, branching and fusion of the newly formed vascular loops, and formation of a new basement membrane. By inhibiting angiogenesis is meant completely or partially inhibiting the formation of such new vasculature.

In certain embodiments, the anti-angiogenic polypeptide is an anti-angiogenic fragment of plasminogen (in particular, angiostatin), an anti-angiogenic fragment of collagen XVIII (endostatin) or a fusion of the two fragments.

Angiostatin is an internal fragment of plasminogen having a molecular weight of 38 or 45 kDa, depending on whether it contains kringles 1-3 or 1-4. In the invention, either can be used, or a molecule including kringles 1-3 and a portion of kringle 4 can be used. Angiostatin can be naturally produced *in vivo* in small amounts by tumor cells, e.g. murine Lewis lung carcinoma cells, by proteolytic cleavage of plasminogen so as to eliminate the N-terminal portion including the signal polypeptide and/or export signal and the preactivation polypeptide and/or export signal, as well as the C-terminal portion following kringle 3 or 4. Mouse and human angiostatin have been purified and sequenced. In preferred embodiments, the gene therapy vectors of this invention encode angiostatin having kringles 1, 2 and 3, or angiostatin having kringles 1, 2, 3 and 4.

In another preferred embodiment, the anti-angiogenic polypeptide is

-7-

endostatin or a biologically active analog or fragment thereof. Endostatin can be naturally produced *in vivo* in small amounts by tumor cells, e.g., murine angiosarcoma cells, by proteolytic cleavage of endogenous collagen XVIII so as to eliminate the N-terminal portion including the signal polypeptide and/or export signal and the preactivation polypeptide and/or export signal, as well as the C-terminal portion following kringle 3 or 4. See Fig.2. Mouse endostatin has been sequenced, and the human molecule (SEQ ID NOs: 17 and 18) forms a portion of collagen 18 (SEQ ID NOs: 19 and 20).

The human molecule position and sequence are apparent from an alignment of the active, Lys-terminated active region of human collagen 18 with murine endostatin, such that the C-terminal lysine residues align, bringing the active endostatin sequences into alignment.

In yet another preferred embodiment, the anti-angiogenic polypeptide is an in-frame fusion of angiostatin or a biologically active analog or fragment thereof and endostatin or a biologically active analog or fragment thereof. Preferably, the angiostatin or biologically active analog or fragment is 5' of the endostatin or biologically active analog or fragment. In certain embodiments, the angiostatin-endostatin fusion proteins exhibit synergistic anti-angiogenic properties.

By fragment is meant some portion of the naturally occurring anti-angiogenic polypeptide. Preferably, the fragment is at least 20 amino acid residues, more preferably at least 50 amino acid residues, and most preferably at least 100 amino acid residues in length. Fragments include chimeric constructs composed of at least a portion of the relevant gene and another molecule. The ability of a candidate fragment to exhibit a biological activity of the anti-angiogenic polypeptide can be assessed by methods known to those skilled in the art, e.g., by its ability to inhibit proliferation of bovine capillary

-8-

cells, or by its ability to inhibit growth of primary tumor cells, e.g., as described herein. See, e.g., Example 9. Also included are fragments containing residues that are not required for biological activity of the fragment or that result from alternative mRNA splicing or alternative protein processing events.

5 Internal or terminal fragments of a polypeptide can be generated by removing one or more nucleotides from one end (for a terminal fragment) or both ends (for an internal fragment) of a nucleic acid which encodes the polypeptide.

In preferred embodiments, the gene therapy vector of this invention is
10 capable of hybridizing to the native anti-angiogenesis polypeptide-encoding regions and has at least about 80%, preferably at least about 90%, and more preferably at least about 95%, sequence identity to the native nucleotide sequences, and encodes a polypeptide which has anti-angiogenic activity; or a biologically active fragment of any of the above nucleotide sequences wherein
15 the encoded polypeptide has anti-angiogenic activity.

The nucleotide sequences of the present invention can be in the form of RNA or DNA, and the nucleotide sequence can be double-stranded or single stranded and, if single stranded, can be the coding strand or non-coding (anti-sense) strand.

20 The coding sequence which encodes the anti-angiogenic polypeptide can be identical to the native coding sequences, or can be a different coding sequence which, as a result of the degeneracy of the genetic code, encodes the same anti-angiogenic polypeptide.

In certain embodiments, the gene therapy vector also has a nucleotide
25 sequence encoding a signal polypeptide and/or export signal (SP) for effecting secretion of the anti-angiogenic polypeptide. Examples of signal polypeptide and/or export signal include plasminogen signal polypeptide and/or export

-9-

signal. Preferably, the signal polypeptide and/or export signal is 5' (i.e., upstream) of the nucleotide sequence encoding the anti-angiogenic polypeptide.

Preferably, the gene therapy vector has a nucleotide sequence encoding a preactivation polypeptide and/or export signal (PAP), which is a small
5 polypeptide and/or export signal which effects folding and secretion of the anti-angiogenic polypeptide *in vivo*. Examples of preactivation polypeptide and/or export signal include plasminogen preactivation polypeptide and/or export signal, described herein, and PAP's of other proteins in the blood clotting cascade.

10 Preferably, the preactivation polypeptide and/or export signal is positioned 5' of the nucleotide sequence encoding the anti-angiogenic polypeptide. In embodiments which have a signal sequence and an anti-angiogenic polypeptide, preferably the preactivation polypeptide and/or export signal is 5' of the nucleotide sequence encoding the anti-angiogenic
15 polypeptide, and 3' of the nucleotide sequence encoding the signal polypeptide and/or export signal.

We have discovered that results obtained using constructs containing a PAP- encoding nucleic acid sequence are far superior to results using constructs lacking a PAP-encoding sequence. Our hypothesis to explain these
20 unexpectedly superior results with PAP is that, during the complex process by which the anti-angiogenic polypeptide is expressed and processed in living cells, the PAP polypeptide and/or export signal facilitates the export of the polypeptide from the cellular Golgi apparatus and/or the endoplasmic reticulum (ER). The corollary is that, absent PAP, a significant portion of the expressed
25 polypeptide remains trapped in the Golgi and/or ER.

The PAP exemplified herein is derived from human plasminogen; this

-10-

PAP is currently preferred. Our discovery that the use of a PAP dramatically improves results leads us to believe that other PAP's would be useful as well, and such others are therefore contemplated for use in the invention. Thus, as used herein, "PAP" refers to a polypeptide and/or export signal which is

5 naturally associated with a eukaryotic (preferably human) protein, the exportation of which is facilitated by its associated PAP. Examples of other human proteins whose Golgi/ER export is PAP-facilitated include other secreted proteins of the blood coagulation cascade, e.g., fibrinogen, prothrombin, Factor VIII, and Factor IX. Other secreted human proteins also

10 are associated with potentially useful PAPs.

It is not essential that the PAP used in the invention be identical in amino acid sequences to a native PAP; it is well-known that polypeptide and/or export signal that facilitate protein secretion or export, e.g., signal polypeptide and/or export signal and PAPs, can vary from the native forms to a certain

15 extent and still retain their function. Therefore, PAPs useful according to the invention preferably have 75% or greater amino acid sequence identity with a native PAP.

In certain embodiments, the gene therapy vector has a nucleotide sequence encoding a tag for identification of the anti-angiogenic polypeptide and/or export signal. In certain embodiments, the tag is 5' of the nucleotide sequence encoding the anti-angiogenic polypeptide; in other embodiments, the tag is 3' of the nucleotide sequence encoding the anti-angiogenic polypeptide.

20 In embodiments in which the anti-angiogenic polypeptide is endostatin or an angiostatin-endostatin fusion, it is preferred that the tag be 5' of the nucleotide sequence encoding endostatin.

25

In certain embodiments the gene therapy vector includes a selectable

-11-

marker, e.g., a Neomycin phosphotransferase gene, or a humanized red-shifted green fluorescent protein.

The invention also includes a cell infected or transfected with a gene therapy vector described herein. Preferably, the cell is an animal cell, more preferably an autologous or allogeneic human cell. The gene therapy vectors described herein can be introduced into a cell, e.g., by transformation, transfection, transduction, infection, or ex vivo injection. They can be targeted to a particular cell type.

Administration of nucleic acid, e.g., a gene therapy vector, can be accomplished by any method which allows the nucleic acid to reach the target cells. These methods include, e.g., injection, deposition, implantation, suppositories, oral ingestion, inhalation, topical administration, or any other method of administration where access to the target cells by the nucleic acid is achieved. Injections can be, e.g., intravenous, intradermal, subcutaneous, intramuscular or intraperitoneal. Implantation includes inserting implantable drug delivery systems, e.g., microspheres, hydrogels, polymeric reservoirs, cholesterol matrices, polymeric systems, e.g., matrix erosion and/or diffusion systems and non-polymeric systems, e.g., compressed, fused or partially fused pellets. Suppositories include glycerin suppositories. Oral ingestion doses can be enterically coated. Inhalation includes administering the nucleic acid with an aerosol in an inhalator, either alone or attached to a carrier that can be absorbed.

In certain embodiments of the invention, administration can be designed so as to result in sequential exposures to the nucleic acid over some time period, e.g., hours, days, weeks, months or years. This can be accomplished by repeated administrations of the nucleic acid, e.g., by one of the methods described above, or alternatively, by a controlled release delivery system in

-12-

which the nucleic acid is delivered to the animal over a prolonged period without repeated administrations. By a controlled release delivery system is meant that total release of the nucleic acid does not occur immediately upon administration, but rather is delayed for some time. Release can occur in bursts or it can occur gradually and continuously. Administration of such a system can be, e.g., by long acting oral dosage forms, bolus injections, transdermal patches or subcutaneous implants. Examples of systems in which release occurs in bursts include, e.g., systems in which the nucleic acid is entrapped in liposomes which are encapsulated in a polymer matrix, the liposomes being sensitive to a specific stimulus, e.g., temperature, pH, light, magnetic field, or a degrading enzyme, and systems in which the nucleic acid agent is encapsulated by an ionically-coated microcapsule with a microcapsule core-degrading enzyme. Examples of systems in which release of the nucleic acid is gradual and continuous include, e.g., erosional systems in which the nucleic acid is contained in a form within a matrix, and diffusional systems in which the nucleic acid permeates at a controlled rate, e.g., through a polymer. Such sustained release systems can be, e.g., in the form of pellets or capsules.

The nucleic acid is administered to the patient in a therapeutically effective amount. By therapeutically effective amount is meant that amount which is capable of at least partially preventing or reversing the disease. A therapeutically effective amount can be determined on an individual basis and will be based, at least in part, on consideration of the patient's size, age, the efficacy of the particular nucleic acid used, the type of delivery system used, the time of administration relative to the onset of disease symptoms, and whether a single, multiple, or controlled release dose regimen is employed. A therapeutically effective amount can be determined by one of ordinary skill in the art employing such factors and using no more than routine experimentation.

-13-

In certain embodiments, a therapeutically effective amount of an anti-angiogenic polypeptide is administered by providing to the animal a nucleic acid encoding the polypeptide and expressing the polypeptide in vivo. Nucleic acids encoding the polypeptide, or mutants thereof, can be administered in any
5 biologically effective carrier, e.g. any formulation or composition capable of effectively delivering the nucleotide sequence for the anti-angiogenic polypeptide to cells in vivo. Approaches include, e.g., insertion of the nucleic acid into viral vectors. Viral vectors can be delivered to the cells, e.g., by infection or transduction using the virus. Viral vectors can also be delivered to
10 the cells, e.g., by physical means, e.g., by electroporation, lipids, cationic lipids, liposomes, DNA gun, $\text{Ca}_3(\text{PO}_4)_2$ precipitation, or delivery of naked DNA. In certain preferred embodiments, the virus is administered by injection, e.g., intramuscular injection, in a dose range of about 10^3 to about 10^{10} infectious particles per injection, more preferably in a dose range of about 10^5 to about
15 10^8 infectious particles per injection. Single or multiple doses can be administered over a given period of time, depending, e.g., upon the disease.

An alternative is insertion of the nucleic acid encoding the anti-angiogenic polypeptide into a bacterial or eukaryotic plasmid. Plasmid DNA can be delivered to cells with the help of, e.g., cationic liposomes (lipofectin™;
20 Life Technologies, Inc., Gaithersburg, MD) or derivatized (e.g., antibody conjugated) polylysine conjugates, gramicidin S, artificial viral envelopes or other such intracellular carriers, as well as direct injection of the gene construct or $\text{Ca}_3(\text{PO}_4)_2$ precipitation carried out in vivo, or by use of a gene gun. The above-described methods are known to those skilled in the art and can be
25 performed without undue experimentation.

Since transfer of the nucleic acid to appropriate target cells represents the critical first step in gene therapy, choice of the particular gene delivery

-14-

system will depend on such factors as the intended target and the route of administration, e.g., locally or systemically. Targets for delivery of the nucleic acid can be, e.g., specific target cells which are diseased. For example, the target can be, e.g., the peritoneal cavity, gastro-intestinal tract, bone marrow
5 cavity, liver, lungs, muscles, vasculature, pericardial cavity, pleural cavity, skin, sub-cutaneous or deep connective tissues, central nervous system, spinal fluid, eye, or specific sites of tumor growth. Administration can be directed to one or more cell types, and to one or more cells within a cell type, so as to be therapeutically effective, by methods known to those skilled in the art. For
10 example, the nucleic acid can be, e.g., coupled to an antibody, to a ligand to a cell surface receptor, or to a toxin component, or can be contained in a particle which is selectively internalized into cells, e.g., liposomes, or a virus where the viral receptor binds specifically to a certain cell type, or a viral particle lacking the viral nucleic acid, or can be administered by local injection.

15 In certain embodiments, the nucleic acid is administered to the patient by introducing *ex vivo* the nucleic acid into cells of the patient, or into syngeneic or allogeneic or xenogeneic cells, and then administering the cells having the nucleic acid to the animal. Any cell type can be used. In certain embodiments, the cells having the introduced nucleic acid are expanded and/or
20 selected after the nucleic acid transfer. The cells having the transferred nucleic acid are subsequently administered to the patient. Preferably, the cells are administered in a dose range of about 1×10^6 to about 1×10^9 cells/dosage/day, and most preferably at about 1×10^7 to about 1×10^8 cells/dosage/day. The cells can be administered by any method which results in delivering the
25 transferred nucleic acid in the cells to the desired target. For example, the cells can be implanted directly into a specific tissue of the patient, or implanted after encapsulation within an artificial polymer matrix. Examples of sites of

-15-

implantation include, e.g., the peritoneal cavity, gastro-intestinal tract, bone marrow cavity, liver, lungs, muscles, vasculature, pericardial cavity, pleural cavity, skin, sub-cutaneous or deep connective tissues, central nervous system, spinal fluid, eye, or specific sites of tumor growth.

5 Systemic delivery can be achieved, e.g., by introducing the nucleic acid into cells which circulate in the peripheral blood of the patient, or which give rise to cells which circulate in the peripheral blood. In certain embodiments, the nucleic acid is introduced into such cells *ex vivo*, and these cells are then administered to the patient, resulting in systemic delivery within the peripheral
10 blood. These cells can be the cells of the patient or allogeneic cells. Preferred cells in which the nucleic acid can be introduced are hematopoietic cells.

 In certain embodiments, other therapy is additionally administered. For example, if the animal is being treated for a tumor, other tumor therapy, e.g., another therapeutic agent, chemotherapy, radiation or surgery, is additionally
15 administered to the patient, either simultaneously or at different times.

 Treating is meant to include, e.g., preventing, treating, reducing the symptoms of, or curing the disease. I.e. treating a tumor includes preventing growth of the tumor, causing shrinkage of the tumor, or preventing development of micro-metastases.

20 Preferably, the recombinant nucleic acid is a gene therapy vector, e.g., as described herein. Preferably, the anti-angiogenic polypeptide is angiostatin, endostatin, an angiostatin-endostatin fusion protein, or biologically active analogs or fragments thereof. In certain embodiments, the angiostatin has kringles 1, 2 and 3; in other embodiments, the angiostatin has kringles 1, 2, 3
25 and 4, and, in some embodiments, kringle 5 of human or murine plasminogen. Angiostatin is described in O'Reilly and Folkman U.S. Patent No. 5,639,725, hereby incorporated by reference. Endostatin is described in O'Reilly and

-16-

Folkman PCT Appln. No. WO 97/15666, published May 1, 1997, hereby incorporated by reference.

In certain embodiments, the recombinant nucleic acid has been introduced *ex vivo* into cells so as to express the anti-angiogenic polypeptide in the cells, and the recombinant nucleic acid is administered to the patient by administering to the patient the cells containing the recombinant nucleic acid. In certain embodiments, the cells are derived from the patient; in other embodiments the cells are allogeneic cells relative to the cells of the patient.

Where cells are infected or transfected *ex vivo* for later infusion into the patient, the cells are preferably hematopoietic cells, but can also be mesenchymal cells, stem cells, epithelial cells (e.g., from the gut), or dendritic cells.

The gene therapy vectors of the invention can be provided in a pharmaceutical composition comprising a therapeutically effective amount of the recombinant nucleic acid together with a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers include, e.g., water, saline, dextrose, glycerol, ethanol, liposomes and lipid emulsions.

The following non-limiting examples further illustrate the present invention.

20 EXAMPLES

Example 1: Construction of Inserts for Gene Therapy Vectors Containing cDNA for Angiostatin, Endostatin or Angiostatin-Endostatin Fusion Proteins

The following genetic constructs are inserted into retroviral gene therapy vectors; the genetic constructs contain human or murine cDNA for angiostatin, endostatin or an angiostatin-endostatin fusion, and DNA encoding a signal

-17-

polypeptide and/or export signal (SP), a tag (FLAG), and, preferably, a preactivation polypeptide and/or export signal (PAP). The constructs are all made using standard genetic engineering techniques, and their insertion into retroviral gene therapy vectors is carried out using known methods. The

5 constructs have the following components:

Murine Constructs

SP-K1-K2-K3-Flag
SP-K1-K2-K3-K4-Flag
SP-K1-K2-K3-K4-K5-Flag
10 SP-PAP-K1-K2-K3-Flag
SP-PAP-K1-K2-K3-K4-Flag (SEQ ID NO: 11 and 12)
SP-Flag-Endo (SEQ ID NO: 13 and 14)
SP-K1-K2-K3-Flag-Endo
SP-K1-K2-K3-K4-Flag-Endo (SEQ ID NO: 15 and 16)
15 SP-PAP-K1-K2-K3-Flag-Endo

Human Constructs

SP-K1-K2-K3
SP-K1-K2-K3-K4
SP-K1-K2-K3-K4-K5
20 SP-PAP-K1-K2-K3
SP-PAP-K1-K2-K3-K4
SP-PAP-K1-K2-K3-K4-K5
SP-Endo
SP-K1-K2-K3-Endo
25 SP-PAP-K1-K2-K3-Endo

Nucleic acid and amino acid sequences for mouse and human angiotensin and endostatin used in these constructs are shown in Figs. 5-7.

Nucleic acid and amino acid sequence of the FLAG peptide:

amino terminus-	ASP	TYR	LYS	ASP	ASP	ASP	ASP	LYS
5'-	GAC	TAC	AAG	GAC	GAC	GAT	GAC	AAG

Human plasminogen derivative constructs

The entire coding region of the human plasminogen cDNA from the start (ATG) to the stop (TAA) codon is 2433bp in size.

This sequence encodes a signal peptide (bp 1-57), a preactivation peptide (bp 58-288), and 5 distinct structural regions known as kringles (K1-K3 from bp 289-1092; K4 from bp 1093-1380; K5 from bp 1381-1740). Please note that although I have given precise bp measurements for kringles K4 and K5, it can be argued that the sequence encoding K4 is between bp 1056-1440 and the sequence encoding K5 is between bp 1362-1680.

A DNA fragment encoding a portion of the human plasminogen protein from bp 1 to 1377 was obtained by PCR of a widely available human liver cDNA library using synthetic DNA oligonucleotides complementary to sequences immediately preceding the signal peptide and immediately following kringle 4. This fragment contains the signal peptide (bp 1-57), the preactivation peptide (bp 58-288), kringles 1 (bp 289-549), 2 (bp 550-804), 3 (bp 805-1092) and 4 (bp 1093-1380). The synthetic oligonucleotides used for this reaction contained engineered recognition sites for the restriction enzymes EcoRI and XhoI. Following the PCR reaction the amplified fragment was cloned into the EcoRI/XhoI sites of BluescriptSK(-) (Stratagene) using standard techniques (Maniatis). Following cloning the integrity of the amplified sequence was verified by sequencing both strands using the Sanger method (Sanger). Various derivatives of the cloned fragment were subsequently constructed using BluescriptSK(-) (Stratagene) as a backbone. A full list of the derivatives are described in Table 1. Briefly, the variations are composed of constructs containing various combinations of kringles with or without the signal and/or preactivation peptide sequences. These derivatives were constructed using both standard techniques as well as PCR and the use of double stranded synthetic oligonucleotides. In all cases the integrity of the start codon, coding sequence and termination codon was verified by double stranded sequencing using the Sanger method.

Murine plasminogen derivative constructs

The coding sequence for murine plasminogen is 2439bp in size and, similar to the human plasminogen cDNA encompasses a sequence encoding signal and preactivation peptides (bp 1-57 and 58-288 respectively) in addition to 5 kringle regions; kringle 1-3 (bp 289-1092), kringle 4 (bp 1093-1380) and kringle 5 (bp 1381-1743). Again, although I have given precise bp measurements for kringles K4 and K5, it can be argued that the sequence encoding K4 is between bp 1056-1440 and the sequence encoding K5 is between bp 1362-1680.

The murine plasminogen cDNA has previously been cloned and was made available to us. Derivatives of murine plasminogen were constructed using sequences derived from bp 1-1743 of the coding sequence. Various combinations of kringle regions with or without signal and preactivation peptide regions were made using BluescriptSK(-) (Stratagene, La Jolla, CA) as the vector backbone. These derivatives were constructed using standard cloning techniques (Maniatis, Molecular cloning; a laboratory manual, second edition, 1989) in combination with PCR utilizing synthetic oligonucleotides using

-19-

Angiostatin function was not altered by adding the FLAG polypeptide and/or export signal to either the N- or C-terminal ends, whereas endostatin was functional only if FLAG was added to its N-terminal end.

Example 2: Construction of Retroviral Gene Therapy Vectors

5 This example illustrates the construction of retroviral gene therapy vectors comprising cDNA for angiostatin, endostatin or angiostatin-endostatin fusion proteins.

 The DNA inserts from Example 1 were inserted into two retroviral vectors. Both vectors were derived from the Murine Stem Cell Virus (MSCV),
10 which is a variant of Moloney Murine Leukemia Virus (MoMLV) having several mutations allowing high, sustained expression in hematopoietic stem cells and their progeny. In both cases, the angiostatin, endostatin, or angiostatin-endostatin fusion DNA inserts were under the transcriptional control of the retroviral left Long Terminal Repeat (LTR). In the first vector,
15 the dominant selectable marker was the Neomycin phosphotransferase gene (NeoR), which confers resistance to G418, and is driven by an internal phosphoglycerate kinase (PGK) promoter. In the second vector, the dominant selectable marker was the humanized, red-shifted green fluorescent protein (EGFP), which is co-translationally expressed by means of an Internal
20 Ribosome Entry Site (IRES) from the Encephalomyocarditis virus (EMCV).

 The retroviral gene therapy vectors were transfected by CaPO_4 precipitation in the transient ecotropic packaging cell-line BOSC 23, Pear et al., *PNAS* 90:8392 (1993). Viral supernatants were collected two days thereafter and filtered through 0.45 mm filters. Filtered viral supernatants were
25 subsequently used to infect GENETIX's stable amphotropic retroviral packaging cell-line AM12 (Genetix Pharmaceuticals, Inc., Cambridge, MA). After another two days, viral supernatants from transduced AM12 were filtered

-20-

and used to infect GENETIX's stable ecotropic retroviral packaging cell-line GP+E86 (Genetix Pharmaceuticals, Inc.). Both transduced AM12 and GP+E86 were then selected in the presence of G418 (in the case of constructs bearing NeoR) or sorted by Fluorescent Activated Cell Sorter (FACS) for EGFP expression. Viral titers were estimated according to standard practice by counting G418 resistant colonies among NIH3T3 cells exposed to diluted virus preparation. Ecotropic viral titers were above 5×10^5 /ml of viral supernatants, only 3-fold lower than "empty" control vectors. No Replication Competent Retrovirus (RCR) was detected in standard assays.

10 Example 3: Transduction of Target Cells Using Retroviral Gene Therapy Vectors

This example illustrates the stability of retroviral gene therapy vector transmission and the lack of toxicity in non-endothelial target cells.

Following 24-hour incubation of confluent viral producer cells in 100 mm plates, viral supernatant was removed and filtered (0.45 μ m filter, Gelman Sciences, Ann Arbor, MI). Viral supernatant, containing 7 μ g/ml polybrene (Sigma, St. Louis, MO), was added to target cells 24 hours after plating the target cells. Fresh medium was added after 4-12 hours, and, after an additional 48 hours, cells were selected for retroviral infection by exposure to medium containing 1 mg/ml G418 (Gibco BRL, Grand Island, NY) or by FACS sorting (FACStar cell sorter, Becton Dickinson, San Jose, CA). The stability of transmission of the retroviral gene therapy vectors described in Example 2 was examined by Southern blot analysis of transduced NIH3T3 cells, using specific probes (EGFP) and restriction enzyme digestion of genomic DNA with SacI, which cuts only once in each LTR. Stable chromosomal integration of intact proviruses of appropriate length was observed with all constructs.

The lack of non-specific toxicity on non-endothelial cells was

-21-

established by using filtered viral supernatants to transduce various tumor cell-types and cell-lines (NIH3T3 cells, K562 cells (ATCC), and human SK-N-AS neuroblastoma cells; Cohen, P.S., *Cancer Research*, 55:2380 (1995).

Transduced cell populations were subsequently selected with G418 or sorted
5 for EGFP expression by FACS. No obvious effects on cell viability, growth or other phenotypical characteristics were detected.

Example 4: Protein Expression of Angiostatin, Endostatin and Angiostatin-Endostatin Fusion Proteins in Cells Transduced with Retroviral Gene Therapy Vectors

10 This example illustrates that recombinant angiostatin, endostatin, and angiostatin-endostatin fusion proteins were readily detected in retrovirally transduced cells and their supernatant, indicating efficient expression and secretion.

MSCV virus based vectors containing sequences encoding murine
15 Kringle 1 (K1), K1K5, K1K2K3, K1K2K3K4, and K1K2K3K4K5 were used to transduce NIH3T3 cells. With regard to the murine recombinant proteins, Western blot analysis of transduced cells and their supernatant was performed by means of a monoclonal antibody that recognizes the FLAG polypeptide and/or export signal. Because this antibody is not mono-specific, significant
20 cross-reactivity with murine proteins was apparent. However, by comparing the pattern obtained with mock cells, it was clear that the antibody revealed an additional band of appropriate size in all transduced cells. Moreover, the recombinant proteins were detected in cell supernatants at levels above 50 ng/ml, using a protein concentration/semi-purification procedure (Centricon
25 columns, Amicon, Beverly, MA). With regard to the human recombinant proteins, no FLAG tag was added, so a monoclonal antibody that recognizes specifically the first three kringles of human plasminogen in its native, non-

-22-

denatured form was used; O'Reilly et al., *Cell* 79:315 (1994). Because of this constraint, Western blot analysis using denaturing gels could not be performed. An ELISA assay was performed which indicated that human recombinant angiostatin was detected at levels likely to be therapeutic according to previous findings in the model of Lewis Lung Carcinoma *Id.*

These results indicate that high levels of recombinant proteins of expected length were expressed in retrovirally transduced cells and were efficiently secreted.

Example 5: In Vivo Anti-Tumor Activity of Cells Transduced with Gene Therapy Vectors Encoding the Angiostatin-Endostatin Fusion Protein

Human SK-N-AS neuroblastoma cells (Cohen, 1995) were transduced with the retroviral gene therapy vector containing the angiostatin-endostatin fusion protein, described in Example 2. These cells (1,000,000) were suspended in 1 mL Dulbecco's phosphate buffered saline and injected into the right mid-quadrant of nude immuno-compromised mice. While no impairment of the *in vitro* growth of transduced cells was observed, a dramatic decrease in tumor growth in nude mice cells following subcutaneous implantation of the transduced cells was evident as compared to "mock virus"-transduced control cells.

Example 6: Ex Vivo Transfer of Retroviral Gene Therapy Vectors Encoding Anti-Angiogenic Polypeptides to Primary Hematopoietic Cells, and Subsequent Transplantation to Recipient Mice

This example illustrates infection of primary hematopoietic cells from

-23-

donor mice with retroviral gene therapy vectors encoding angiostatin, endostatin, or an angiostatin-endostatin fusion protein, and a selectable GFP marker, and subsequent transplantation of the transduced hematopoietic cells into recipient mice.

- 5 Femoral bone marrow cells are harvested from male donor C57BL6/J-Ly5.1 mice (Jackson Labs, Bar Harbor, ME), intravenously injected four days previously with 150 mg/kg of 5-fluorouracil (5-FU). Bone marrow cells are cultured for two days in medium composed of DMEM, 15% fetal calf serum, 10 ng/ml human IL-6, 6 ng/ml murine IL-3 and 100 ng/ml murine Steel factor
10 prior to two days of culture atop a confluent monolayer of irradiated (1,500 cGy, ^{137}Cs γ -irradiation) viral producer cells in the above medium including 6 ug/ml of prolamine sulfate. The viral producer cells are transfected with a retroviral gene therapy vector, as described above. Upon completion of the co-culture infection protocol, recovered non-adherent cells are cultured for an
15 additional 48 hours to allow for expression of the transferred GFP gene. Retrovirally transduced cells expressing the transferred GFP gene are subsequently identified and selected for, using a FACStar+ cell sorter (Becton Dickinson, San Jose, CA). The GFP+ cells are intravenously injected into congenic female C57BL6/J-Ly5.2 recipient mice (National Cancer Institute,
20 Washington, DC) previously given 950 cGy (83cGy/min, ^{137}Cs γ -rays) of whole body irradiation. In each case, a small fraction of GFP+ sorted cells is used for day 12 CFU-S and *in vitro* clonogenic progenitor assays to assess the efficiency of the infection and selection procedures on these more mature cell types.

25 Example 7: Engraftment of Recipient Mice with Donor-Derived Hematopoietic Cells

This example illustrates engraftment of the recipient mice with the

-24-

donor-derived transfected hematopoietic cells from Example 6.

The donor and recipient mice are phenotypically distinguishable on the basis of Y chromosome specific sequences, as well as on the basis of allelic differences at the murine CD45 cell surface antigen locus. Male donor mice are homozygous for the CD45.2 allele, while female recipient mice are homozygous for CD45.1. The engraftment of recipient mice with donor-derived (CD45.2+) cells is assessed at both short (5 weeks) and long (34 months) time points post-transplant by flow cytometric analysis of peripheral blood samples stained with a phycoerythrin labeled antibody specific for the CD45.2 antigen (Pharmingen, San Diego, CA). The results indicate that engraftment occurs.

Example 8: Proviral Marking and GFP Expression in Recipient Mice

This example illustrates the presence of recombinant provirus and expression of the transferred GFP gene in the recipient mice from Example 6.

The level of proviral marking in reconstituted animals is initially determined by Southern blot and semi-quantitative PCR analysis of DNA obtained from peripheral blood leukocytes. The large majority of donor-derived (CD45.2+) cells in recipient mice contain a minimum of one copy of recombinant provirus. In addition, flow cytometric analysis of peripheral blood leukocytes is performed to ascertain the proportion of cells expressing the transferred GFP cDNA. Because the GFP and angiogenic inhibitor protein cDNAs are both driven from the same regulatory sequences, due to the inclusion of an internal ribosomal entry site (IRES) element, the analysis of GFP expression in the peripheral blood provides an indirect measurement of the levels of anti-angiogenic protein being expressed. The results indicate expression of the transferred genes.

Example 9: Anti-Angiogenic Polypeptide Expression in Recipient Mice

This example illustrates the presence of anti-angiogenic polypeptide in the sera of the recipient mice from Example 6, using both physical and functional assays.

5 Serum obtained from the transplanted animals described in Example 6 is used for ELISA using an antibody specific for the synthetic FLAG epitope (IBI, Eastman Kodak, New Haven, CT) and compared against known standards of purified protein. Results indicate the presence of the anti-angiogenic polypeptide in the serum.

10 To determine whether a functional anti-angiogenesis polypeptide is present in the circulation, sera from transplanted animals is tested for its ability to inhibit the proliferation of bovine capillary cells *in vitro*; O'Reilly (1994). Briefly, cells are plated in 24 well dishes at 25,000 cells/ml and maintained in DMEM with 5% bovine calf serum for 24 hours. The medium is then replaced
15 with fresh medium containing various dilutions of the test serum. After 20 minutes of incubation, fresh medium including b-FGF (final concentration 1 ng/ml) is added and the cells are cultured for 72 hours. Cells are then dispersed using trypsin and the cell number determined by Coulter counter. Results indicate that functional anti-angiogenic polypeptide is present in the sera of the
20 recipient mice.

 In addition, the ability of circulating anti-angiogenic polypeptide to inhibit the growth of primary tumor cells is assessed. Transplanted mice are subcutaneously injected with one million Lewis lung carcinoma (LLC) cells (O'Reilly, (1994)) at the proximal midline of their dorsal skin. The mice are
25 closely monitored for survival, tumor size and growth, and overall health. Results indicate that the anti-angiogenic polypeptides from the sera of the

-26-

recipient mice inhibit growth of the LLC tumor cells.

Finally, upon sacrifice of the transplanted recipient mice, blood, spleen, thymus and bone marrow are harvested and analyzed for the presence of proviral DNA by Southern analysis as well as expression of the transferred

5 GFP and anti-angiogenic polypeptide cDNAs by flow cytometry and ELISA. Moreover, a portion of bone marrow cells is re-transplanted into secondary recipients to generate individual day 12 spleen colonies, as well as plated in methylcellulose to assess *in vitro* clonogenic progenitors. Individual clones are analyzed for proviral DNA by PCR or Southern blot, and for gene expression

10 by flow cytometry and ELISA. Results of these tests also indicate the presence of proviral DNA and expression of the anti-angiogenic polypeptides and marker proteins.

Example 10: Evaluating the Efficacy of Retroviral Gene Therapy Vectors
Encoding Anti-Angiogenic Polypeptides on Various Human
Cancers Implanted in SCID Mice Using Ex Vivo Gene Therapy

15

This example illustrates a method for rapidly screening various forms of human cancer to determine susceptibility to treatment by the systemic delivery of anti-angiogenic polypeptides.

The methods for gene transfer, assessment of proviral marking and

20 assessment of transferred gene expression as described in Examples 3 through 9 are repeated using immuno-deficient SCID mice, with the following exceptions. Since SCID mice are more sensitive to γ -irradiation than C57BL6/J mice, the female SCID recipients receive a lower dose of 400cGy of whole body irradiation in contrast to the 950cGy required for C57BL6/J. In

25 addition, since the SCID mice do not possess allelic differences at the CD45 cell surface antigen locus, donor and recipient cells are phenotypically distinguished on the basis of Y chromosome specific sequences using Southern

-27-

blot analysis.

Bone marrow from male donor SCID mice is infected, selected for on the basis of expression of the transferred GFP marker cDNA, and transplanted into irradiated female SCID recipients. Engraftment with provirally marked
5 cells and expression of the transferred genes is demonstrated. The mice are then separately implanted with a variety of human tumor cell types, e.g., breast adenocarcinoma, lung squamous cell carcinoma, and brain glioblastoma. In each case, the ability of the anti-angiogenic polypeptides to inhibit the growth of the various human tumor cell types is monitored and quantified.

10 Example 11: Evaluating the Efficacy of Retroviral Gene Therapy Vectors Encoding Anti-Angiogenic Polypeptides for Treatment of Ovarian Cancer Using In Vivo Gene Therapy

This example illustrates the feasibility of using retroviral gene therapy vectors encoding anti-angiogenic polypeptides to achieve efficient gene transfer
15 to established tumors *in vivo* using a well-established murine model of human ovarian cancer. Following injections, mice are closely monitored for tumor growth and survival.

Eight to ten week old nude mice (Jackson Labs, Bar Harbor) are injected intra-peritoneally with 1×10^7 PA-1 cells, an ovarian cancer cell-line (ATCC),
20 and followed until palpable tumors are identified. Viral supernatant for *in vivo* injection is prepared as follows: Viral producer cells are grown to confluence in DMEM with 10% bovine calf serum, and the medium is then changed. After 24 hours of incubation, the viral conditioned supernatant is filtered through a 0.45 μ m low protein binding filter, protamine sulfate is added to a final
25 concentration of 6 μ g/ml, the solution is aliquoted into 2 ml volumes, and frozen at -80°C . Recipient mice receive three intraperitoneal injections of viral supernatant (2 mls per injection) in addition to the polycation, over a period of

-28-

36 hours. Control mice are injected with medium collected from confluent dishes of NIH3T3 cells. Following injection of the viral conditioned supernatant, the mice are analyzed for survival as well as tumor growth over time as compared to mock injected controls. Results indicate that treatment of
5 the ovarian cancer occurs. At death, the tumors are removed, weighed, and the cells dissociated for DNA extraction for Southern blot analysis to detect recombinant provirus.

Those skilled in the art will be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the
10 invention described herein. These and all other equivalents are intended to be encompassed by the following claims.

In other embodiments, the invention provides methods and compositions for treating diseases and processes that are mediated by angiogenesis including, but not limited to, hemangioma, solid tumors, leukemia, metastasis,
15 telangiectasia, psoriasis, scleroderma, pyogenic granuloma, myocardial angiogenesis, plaque neovascularization, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, corneal diseases, rubeosis, neovascular glaucoma, diabetic retinopathy, retrolental fibroplasia, arthritis, diabetic neovascularization, macular degeneration, wound
20 healing, peptic ulcer, *Helicobacter* related diseases, fractures, keloids, vasculogenesis, hematopoiesis, ovulation, menstruation, placentation, and cat scratch fever.

What is claimed is:

-29-

CLAIMS

1. Use of a nucleic acid molecule which expresses an anti-angiogenic polypeptide selected from the group consisting of human angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fragments thereof in the preparation of a medicament for inhibiting tumor growth in a human patient harboring a solid tumor, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth.

2. Use of a nucleic acid molecule which expresses an anti-angiogenic polypeptide of the formula A-B, wherein
A and B are polypeptide and/or export signal joined by a polypeptide and/or export signal bond;

15 A contains at least amino acids and comprises kringles 1, 2, and 3 of human or murine angiostatin; and

B contains at least amino acids and includes at least 75% of the amino acid sequence of human or murine endostatin in the preparation of a medicament for inhibiting tumor growth in a human patient harboring a solid tumor, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth.

3. The use of claim 2, wherein A further comprises kringle region 4 of

-30-

human or murine angiostatin.

4. The use of claim 2 or claim 3, wherein A further comprises kringle 5 of human or murine plasminogen.

5. The use of claim 1 or claim 2, wherein said nucleic acid molecule constitutes a portion of a viral vector.

6. The use of claim 1 or claim 2, wherein said nucleic acid molecule constitutes a portion of a plasmid.

7. The use of claim 6, wherein said plasmid is carried in a cell-free carrier so that the plasmid transfects living cells of the patient following plasmid administration, causing expression of the anti-angiogenesis polypeptide and/or export signal in the patient such that angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells is inhibited, causing inhibition of tumor growth.

8. The use of claim 6, wherein said plasmid has been transfected into animal cells *ex vivo*, wherein said animal cells express the anti-angiogenesis polypeptide to inhibit tumor-associated angiogenesis and tumor growth.

9. The use of claim 5, wherein said viral vector is carried in a cell-free carrier, so that the viral vector is incorporated into living cells of the patient following viral vector administration, causing expression of the anti-

-31-

angiogenesis polypeptide in the patient such that angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells is inhibited, causing inhibition of tumor growth.

5 10. The use of claim 5, wherein animal cells are infected with said viral vector *ex vivo* and then administered to said patient, wherein said animal cells express the anti-angiogenesis polypeptide to inhibit tumor-associated angiogenesis and tumor growth.

11. The use of claim 8, wherein said animal cells are human cells.

10 12. The use of claim 11, wherein said human cells are autologous.

13. The use of claim 11, wherein said human cells are allogeneic.

14. The use of claim 10, wherein said animal cells are human cells.

15. The use of claim 14, wherein said human cells are autologous.

16. The use of claim 14, wherein said human cells are allogeneic.

15 17. The use of claim 5, wherein said viral vector is a retroviral vector.

18. The use of claim 5, wherein said viral vector is a non-retroviral vector selected from the group consisting of adenoviral, adeno-associated, herpes, Simliki Forest virus, and poxvirus vectors.

-32-

19. The use of claim 17, wherein said retroviral vector is Murine Stem Cell Virus or a lentivirus.

20. The use of claim 1, wherein said angiostatin comprises kringles 1, 2 and 3.

5 21. The use of claim 20, wherein said angiostatin further comprises kringle 4.

22. The use of claim 1, wherein said anti-angiogenic polypeptide is a fusion of angiostatin or a biologically active fragment thereof and endostatin or a biologically active fragment thereof.

10 23. The use of claim 1, wherein said nucleic acid molecule includes a nucleotide sequence encoding a signal polypeptide and/or export signal for effecting secretion of said anti-angiogenesis polypeptide.

24. The use of claim 23, wherein said signal polypeptide and/or export signal is plasminogen signal polypeptide and/or export signal.

15 25. The use of claim 1, wherein said nucleic acid molecule includes a nucleotide sequence encoding a preactivation polypeptide and/or export signal for effecting Golgi and/or ER export of the anti-angiogenic polypeptide..

26. The use of claim 25, wherein said preactivation polypeptide and/or export signal is a preactivation polypeptide and/or export signal of a human
20 protein of the blood coagulation cascade.

27. The use of claim 26, wherein said preactivation polypeptide and/or export signal is human plasminogen preactivation polypeptide and/or export signal.

28. The method of claim 25, wherein the preactivation encoding
5 sequence is positioned between a signal-encoding sequence and the sequence encoding the anti-angiogenic polypeptide and/or export signal.

29. The use of claim 1, wherein said nucleic acid molecule includes a nucleotide sequence encoding a tag for identification of said anti-angiogenic polypeptide.

10 30. The method of claim 27, wherein said tag is a Flag tag polypeptide and/or export signal.

31. A viral gene therapy vector comprising a nucleic acid molecule which encodes an anti-angiogenic polypeptide selected from the group consisting of human angiostatin, murine angiostatin, human endostatin, murine
15 endostatin, and angiogenesis-inhibiting fusions and fragments thereof, wherein said viral vector is sufficiently attenuated for use in human gene therapy.

32. A human cell infected with the vector of claim 31.

33. Use of a nucleic acid molecule which expresses in said patient an anti-angiogenic polypeptide selected from the group consisting of human
20 angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fusions and fragments thereof, in the preparation of a

medicament for treating a human patient suffering from diabetic retinopathy, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the retina.




FIG. 1




Fig. 2





FIG. 3



17:1

FIG. 4

5/8




Fig. 5

[illegible]

end mouse endostatin coding sequence

Fig 6

Mouse Plasmid no gen cDNA




FIG. 7 (Sheet 1 of 2)

Tue, Oct 21, 1997									
Code: Universal									
Mou Plasm cDNA									
Page 3									
2010	2020	2030	2040	2050	2060	2070	2080	2090	
CATTGCCCTGCTGAACTAAGCCGCCAGCCACCATCAGGATAAAGTCATTCCAGCTGTCTGCCATCTCCAAATTACATGGTTGCTGA									
IleAlaLeuLeuLysLeuSerArgProAlaThrIleThrAspLysValIleProAlaCysLeuProSerProAsnTyrMetValAlaAsp									
2100									
CCGGACAATA									
ArgThrIle									
2110	2120	2130	2140	2150	2160	2170	2180	2190	
TGTTCATCTACCCGGCTGGGGAGAGACTCAAGGGACTTTCGGTGCCGGTCTGCTCAAGGAGGCTCAGCTGCCCTCTGATTGAGAACAAGGTG									
CysTyrIleThrGlyTrpGlyGluThrGlnGlyThrPheGlyAlaGlyArgLeuLysGluAlaGlnLeuProValIleGluAsnLysVal									
2200									
TGCAACCGCG									
CysAsnArgVal									
2210	2220	2230	2240	2250	2260	2270	2280	2290	
TCGAGTATCTGAACAACAGACTCAATCCACGGAGCTCTGTGCCGGGCAACTGGCTGGTGGCGTCGACAGCTGCCAAGGCGACAGTGGAG									
GluTyrLeuAsnAsnArgValLysSerThrGluLeuCysAlaGlyGlnLeuAlaGlyGlyValAspSerCysGlnGlyAspSerGlyGly									
2300									
GACCTCTGGT									
ProLeuVal									
2310	2320	2330	2340	2350	2360	2370	2380	2390	
TTGCTTCGAGAAGGACAAGTACATTTTACAAGGAGTCACTTCTTGGGGTCTTGGCTGTGCTCGCCCCAATAAGCCTGGTGTCTACGTTCTG									
CysPheGluLysAspLysTyrIleLeuGlnGlyValThrSerTrpGlyLeuGlyCysAlaArgProAsnLysProGlyValTyrValArg									
2400									
TGCTCACGG									
ValSerArg									
2410	2420	2430	2440	2450	2460	2470	2480	2490	
TTTGTTCATTGGATTGAAAGGGAGATGAGGAATAACGACTAGGTGGAAGGCCGAGCAAAACCTCTGCTTACTAAAGCTTACTGAATATG									
PheValAspTrpIleGluArgGluMetArgAsnAsnEndLeuGlyGlyArgProSerLysThrSerAlaTyrEndSerLeuLeuAsnMet									
2500									
GGGAGAGGCC									
GlyArgGlyLeu									
2510	2520	2530	2540	2550	2560	2570	2580	2590	
TTAGGGTGTTTGGAAAACTGACAGTAATCAAACTGGGACACTACACTGAACACAGCTTCCTGTGCCCCCTCAGCCCCCTCCCTTTT									
ArgValPheGlyLysThrAspSerAsnGlnThrGlyThrLeuHisEndThrThrAlaSerCysArgProSerAlaProProLeuPheLeu									
2600									
TGTATTATTG									
TyrTyrCys									
2610	2620	2630	2640	2650	2660	2670	2680	2690	
TGGGTAAAAATTTCTGTCTGTGGACTTCTGGATTTGTGACAATAGACCATCACTGTGTGACCTTTGTGAAAAATAAACTCGATACTT									
GlyEndAsnPheProValCysGlyLeuLeuAspPheValThrIleAspHisHisCysCysAspLeuCysEndLysEndThrArgTyrLeu									
2700									
ACTTTG									
Leu									

FIG. 7 (Sheet 2 of 2)

SEQUENCE LISTING

<110> Genetix Pharmaceuticals, Inc.

<120> ANTI-ANGIOGENIC GENE THERAPY VECTORS AND
THEIR USE IN TREATING ANGIOGENESIS-RELATED DISEASES

<130> 50033/002W01

<140> PCT/US98/xxxxx

<141> 1998-11-20

<150> 08/975,424

<151> 1997-11-20

<160> 20

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2433

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)...(2433)

<400> 1

atg gaa cat aag gaa gtg gtt ctt cta ctt ctt tta ttt ctg aaa tca	48
Met Glu His Lys Glu Val Val Leu Leu Leu Leu Phe Leu Lys Ser	
1 5 10 15	
ggg caa gga gag cct ctg gat gac tat gtg aat acc cag ggg gct tca	96
Gly Gln Gly Glu Pro Leu Asp Asp Tyr Val Asn Thr Gln Gly Ala Ser	
20 25 30	
ctg ttc agt gtc act aag aag cag ctg gga gca gga agt ata gaa gaa	144
Leu Phe Ser Val Thr Lys Lys Gln Leu Gly Ala Gly Ser Ile Glu Glu	
35 40 45	
tgt gca gca aaa tgt gag gag gac gaa gaa ttc acc tgc agg gca ttc	192
Cys Ala Ala Lys Cys Glu Glu Asp Glu Glu Phe Thr Cys Arg Ala Phe	
50 55 60	
caa tat cac agt aaa gag caa caa tgt gtg ata atg gct gaa aac agg	240
Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Arg	
65 70 75 80	
aag tcc tcc ata atc att agg atg aga gat gta gtt tta ttt gaa aag	288
Lys Ser Ser Ile Ile Arg Met Arg Asp Val Val Leu Phe Glu Lys	
85 90 95	

aaa gtg tat ctc tca gag tgc aag act ggg aat gga aag aac tac aga Lys Val Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg 100 105 110	336
ggg acg atg tcc aaa aca aaa aat ggc atc acc tgt caa aaa tgg agt Gly Thr Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser 115 120 125	384
tcc act tct ccc cac aga cct aga ttc tca cct gct aca cac ccc tca Ser Thr Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser 130 135 140	432
gag gga ctg gag gag aac tac tgc agg aat cca gac aac gat ccg cag Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln 145 150 155 160	480
ggg ccc tgg tgc tat act act gat cca gaa aag aga tat gac tac tgc Gly Pro Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys 165 170 175	528
gac att ctt gag tgt gaa gag gaa tgt atg cat tgc agt gga gaa aac Asp Ile Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn 180 185 190	576
tat gac ggc aaa att tcc aag acc atg tct gga ctg gaa tgc cag gcc Tyr Asp Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala 195 200 205	624
tgg gac tct cag agc cca cac gct cat gga tac att cct tcc aaa ttt Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe 210 215 220	672
cca aac aag aac ctg aag aag aat tac tgt cgt aac ccc gat agg gag Pro Asn Lys Asn Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu 225 230 235 240	720
ctg cgg cct tgg tgt ttc acc acc gac ccc aac aag cgc tgg gaa ctt Leu Arg Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu 245 250 255	768
tgc gac atc ccc cgc tgc aca aca cct cca cca tct tct ggt ccc acc Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Ser Gly Pro Thr 260 265 270	816
tac cag tgt ctg aag gga aca ggt gaa aac tat cgc ggg aat gtg gct Tyr Gln Cys Leu Lys Gly Thr Gly Glu Asn Tyr Arg Gly Asn Val Ala 275 280 285	864
gtt acc gtt tcc ggg cac acc tgt cag cac tgg agt gca cag acc cct Val Thr Val Ser Gly His Thr Cys Gln His Trp Ser Ala Gln Thr Pro 290 295 300	912
cac aca cat aac agg aca cca gaa aac ttc ccc tgc aaa aat ttg gat His Thr His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Asp 305 310 315 320	960

gaa aac tac tgc cgc aat cct gac gga aaa agg gcc cca tgg tgc cat	1008
Glu Asn Tyr Cys Arg Asn Pro Asp Gly Lys Arg Ala Pro Trp Cys His	
325 330 335	
aca acc aac agc caa gtg cgg tgg gag tac tgt aag ata ccg tcc tgt	1056
Thr Thr Asn Ser Gln Val Arg Trp Glu Tyr Cys Lys Ile Pro Ser Cys	
340 345 350	
gac tcc tcc cca gta tcc acg gaa caa ttg gct ccc aca gca cca cct	1104
Asp Ser Ser Pro Val Ser Thr Glu Gln Leu Ala Pro Thr Ala Pro Pro	
355 360 365	
gag cta acc cct gtg gtc cag gac tgc tac cat ggt gat gga cag agc	1152
Glu Leu Thr Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Gln Ser	
370 375 380	
tac cga ggc aca tcc tcc acc acc acc aca gga aag aag tgt cag tct	1200
Tyr Arg Gly Thr Ser Ser Thr Thr Thr Thr Gly Lys Lys Cys Gln Ser	
385 390 395 400	
tgg tca tct atg aca cca cac cgg cac cag aag acc cca gaa aac tac	1248
Trp Ser Ser Met Thr Pro His Arg His Gln Lys Thr Pro Glu Asn Tyr	
405 410 415	
cca aat gct ggc ctg aca atg aac tac tgc agg aat cca gat gcc gat	1296
Pro Asn Ala Gly Leu Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp	
420 425 430	
aaa ggc ccc tgg tgt ttt acc aca gac ccc agc gtc agg tgg gag tac	1344
Lys Gly Pro Trp Cys Phe Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr	
435 440 445	
tgc aac ctg aaa aaa tgc tca gga aca gaa gcg agt gtt gta gca cct	1392
Cys Asn Leu Lys Lys Cys Ser Gly Thr Glu Ala Ser Val Val Ala Pro	
450 455 460	
ccg cct gtt gtc ctg ctt cca gat gta gag act cct tcc gaa gaa gac	1440
Pro Pro Val Val Leu Leu Pro Asp Val Glu Thr Pro Ser Glu Glu Asp	
465 470 475 480	
tgt atg ttt ggg aat ggg aaa gga tac cga ggc aag agg gcg acc act	1488
Cys Met Phe Gly Asn Gly Lys Gly Tyr Arg Gly Lys Arg Ala Thr Thr	
485 490 495	
gtt act ggg acg cca tgc cag gac tgg gct gcc cag gag ccc cat aga	1536
Val Thr Gly Thr Pro Cys Gln Asp Trp Ala Ala Gln Glu Pro His Arg	
500 505 510	
cac agc att ttc act cca gag aca aat cca cgg gcg ggt ctg gaa aaa	1584
His Ser Ile Phe Thr Pro Glu Thr Asn Pro Arg Ala Gly Leu Glu Lys	
515 520 525	
aat tac tgc cgt aac cct gat ggt gat gta ggt ggt ccc tgg tgc tac	1632
Asn Tyr Cys Arg Asn Pro Asp Gly Asp Val Gly Gly Pro Trp Cys Tyr	

530	535	540	
acg aca aat cca aga aaa ctt tac gac tac tgt gat gtc cct cag tgt			1680
Thr Thr Asn Pro Arg Lys Leu Tyr Asp Tyr Cys Asp Val Pro Gln Cys			
545	550	555	560
gcg gcc cct tca ttt gat tgt ggg aag cct caa gtg gag ccg aag aaa			1728
Ala Ala Pro Ser Phe Asp Cys Gly Lys Pro Gln Val Glu Pro Lys Lys			
565	570	575	
tgt cct gga agg gtt gtg ggg ggg tgt gtg gcc cac cca cat tcc tgg			1776
Cys Pro Gly Arg Val Val Gly Gly Cys Val Ala His Pro His Ser Trp			
580	585	590	
ccc tgg caa gtc agt ctt aga aca agg ttt gga atg cac ttc tgt gga			1824
Pro Trp Gln Val Ser Leu Arg Thr Arg Phe Gly Met His Phe Cys Gly			
595	600	605	
ggc acc ttg ata tcc cca gag tgg gtg ttg act gct gcc cac tgc ttg			1872
Gly Thr Leu Ile Ser Pro Glu Trp Val Leu Thr Ala Ala His Cys Leu			
610	615	620	
gag aag tcc cca agg cct tca tcc tac aag gtc atc ctg ggt gca cac			1920
Glu Lys Ser Pro Arg Pro Ser Ser Tyr Lys Val Ile Leu Gly Ala His			
625	630	635	640
caa gaa gtg aat ctc gaa ccg cat gtt cag gaa ata gaa gtg tct agg			1968
Gln Glu Val Asn Leu Glu Pro His Val Gln Glu Ile Glu Val Ser Arg			
645	650	655	
ctg ttc ttg gag ccc aca cga aaa gat att gcc ttg cta aag cta agc			2016
Leu Phe Leu Glu Pro Thr Arg Lys Asp Ile Ala Leu Leu Lys Leu Ser			
660	665	670	
agt cct gcc gtc atc act gac aaa gta atc cca gct tgt ctg cca tcc			2064
Ser Pro Ala Val Ile Thr Asp Lys Val Ile Pro Ala Cys Leu Pro Ser			
675	680	685	
cca aat tat gtg gtc gct gac cgg acc gaa tgt ttc atc act ggc tgg			2112
Pro Asn Tyr Val Val Ala Asp Arg Thr Glu Cys Phe Ile Thr Gly Trp			
690	695	700	
gga gaa acc caa ggt act ttt gga gct ggc ctt ctc aag gaa gcc cag			2160
Gly Glu Thr Gln Gly Thr Phe Gly Ala Gly Leu Leu Lys Glu Ala Gln			
705	710	715	720
ctc cct gtg att gag aat aaa gtg tgc aat cgc tat gag ttt ctg aat			2208
Leu Pro Val Ile Glu Asn Lys Val Cys Asn Arg Tyr Glu Phe Leu Asn			
725	730	735	
gga aga gtc caa tcc acc gaa ctc tgt gct ggg cat ttg gcc gga ggc			2256
Gly Arg Val Gln Ser Thr Glu Leu Cys Ala Gly His Leu Ala Gly Gly			
740	745	750	
act gac agt tgc cag ggt gac agt gga ggt cct ctg gtt tgc ttc gag			2304

Thr Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Phe Glu
 755 760 765
 aag gac aaa tac att tta caa gga gtc act tct tgg ggt ctt ggc tgt 2352
 Lys Asp Lys Tyr Ile Leu Gln Gly Val Thr Ser Trp Gly Leu Gly Cys
 770 775 780
 gca cgc ccc aat aag cct ggt gtc tat gtt cgt gtt tca agg ttt gtt 2400
 Ala Arg Pro Asn Lys Pro Gly Val Tyr Val Arg Val Ser Arg Phe Val
 785 790 795 800
 act tgg att gag gga gtg atg aga aat aat taa 2433
 Thr Trp Ile Glu Gly Val Met Arg Asn *
 805 810

<210> 2
 <211> 810
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Glu His Lys Glu Val Val Leu Leu Leu Leu Leu Phe Leu Lys Ser
 1 5 10 15
 Gly Gln Gly Glu Pro Leu Asp Asp Tyr Val Asn Thr Gln Gly Ala Ser
 20 25 30
 Leu Phe Ser Val Thr Lys Lys Gln Leu Gly Ala Gly Ser Ile Glu Glu
 35 40 45
 Cys Ala Ala Lys Cys Glu Glu Asp Glu Glu Phe Thr Cys Arg Ala Phe
 50 55 60
 Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Arg
 65 70 75 80
 Lys Ser Ser Ile Ile Ile Arg Met Arg Asp Val Val Leu Phe Glu Lys
 85 90 95
 Lys Val Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg
 100 105 110
 Gly Thr Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser
 115 120 125
 Ser Thr Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser
 130 135 140
 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln
 145 150 155 160
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys
 165 170 175
 Asp Ile Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn
 180 185 190
 Tyr Asp Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala
 195 200 205
 Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe
 210 215 220
 Pro Asn Lys Asn Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu
 225 230 235 240
 Leu Arg Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu
 245 250 255
 Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Ser Gly Pro Thr

6

Gly Glu Thr Gln Gly Thr Phe Gly Ala Gly Leu Leu Lys Glu Ala Gln
 705 710 715 720
 Leu Pro Val Ile Glu Asn Lys Val Cys Asn Arg Tyr Glu Phe Leu Asn
 725 730 735
 Gly Arg Val Gln Ser Thr Glu Leu Cys Ala Gly His Leu Ala Gly Gly
 740 745 750
 Thr Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Phe Glu
 755 760 765
 Lys Asp Lys Tyr Ile Leu Gln Gly Val Thr Ser Trp Gly Leu Gly Cys
 770 775 780
 Ala Arg Pro Asn Lys Pro Gly Val Tyr Val Arg Val Ser Arg Phe Val
 785 790 795 800
 Thr Trp Ile Glu Gly Val Met Arg Asn Asn
 805 810

<210> 3

<211> 2696

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1) ... (2436)

<400> 3

atg gac cat aag gaa gta atc ctt ctg ttt ctc ttg ctt ctg aaa cca	48
Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro	
1 5 10 15	
gga caa ggg gac tcg ctg gat ggc tac ata agc aca caa ggg gct tca	96
Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser	
20 25 30	
ctg ttc agt ctc acc aag aag cag ctc gca gca gga ggt gtc tcg gac	144
Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp	
35 40 45	
tgt ttg gcc aaa tgt gaa ggg gaa aca gac ttt gtc tgc agg tca ttc	192
Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe	
50 55 60	
cag tac cac agc aaa gag cag caa tgc gtg atc atg gcg gag aac agc	240
Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Ser	
65 70 75 80	
aag act tcc tcc atc atc cgg atg aga gac gtc atc tta ttc gaa aag	288
Lys Thr Ser Ser Ile Ile Arg Met Arg Asp Val Ile Leu Phe Glu Lys	
85 90 95	
aga gtg tat ctg tca gaa tgt aag acc ggc atc ggc aac ggc tac aga	336
Arg Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg	
100 105 110	
gga acc atg tcc agg aca aag agt ggt gtt gcc tgt caa aag tgg ggt	384
Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly	

115	120	125	
gcc acg ttc ccc cac gta ccc aac tac tct ccc agt aca cat ccc aat			432
Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn			
130	135	140	
gag gga cta gaa gag aac tac tgt agg aac cca gac aat gat gaa caa			480
Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln			
145	150	155	160
ggg cct tgg tgc tac act aca gat ccg gac aag aga tat gac tac tgc			528
Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys			
165	170	175	
aac att cct gaa tgt gaa gag gaa tgc atg tac tgc agt gga gaa aag			576
Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys			
180	185	190	
tat gag ggc aaa atc tcc aag acc atg tct gga ctt gac tgc cag gcc			624
Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala			
195	200	205	
tgg gat tct cag agc cca cat gct cat gga tac atc cct gcc aaa ttt			672
Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe			
210	215	220	
cca agc aag aac ctg aag atg aat tat tgc cac aac cct gac ggg gag			720
Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu			
225	230	235	240
cca agg ccc tgg tgc ttc aca aca gac ccc acc aaa cgc tgg gaa tac			768
Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr			
245	250	255	
tgt gac atc ccc cgc tgc aca aca ccc ccg ccc cca ccc agc cca acc			816
Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr			
260	265	270	
tac caa tgt ctg aaa gga aga ggt gaa aat tac cga ggg acc gtg tct			864
Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser			
275	280	285	
gtc acc gtg tct ggg aaa acc tgt cag cgc tgg agt gag caa acc cct			912
Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro			
290	295	300	
cat agg cac aac agg aca cca gaa aat ttc ccc tgc aaa aat ctg gaa			960
His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu			
305	310	315	320
gag aac tac tgc ccg aac cca gat gga gaa act gct ccc tgg tgc tat			1008
Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr			
325	330	335	
acc act gac agc cag ctg agg tgg gag tac tgt gag att cca tcc tgc			1056

Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys	
340 345 350	
gag tcc tca gca tca cca gac cag tca gat tcc tca gtt cca cca gag	1104
Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu	
355 360 365	
gag caa aca cct gtg gtc cag gaa tgc tac cag agc gat ggg cag agc	1152
Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser	
370 375 380	
tat cgg ggt aca tcg tcc act acc atc aca ggg aag aag tgc cag tcc	1200
Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser	
385 390 395 400	
tgg gca gct atg ttt cca cac agg cat tcg aag acc cca gag aac ttc	1248
Trp Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe	
405 410 415	
cca gat gct ggc ttg gag atg aac tac tgc agg aac ccg gat ggt gac	1296
Pro Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp	
420 425 430	
aag ggc cct tgg tgc tac acc act gac ccg agc gtc agg tgg gaa tac	1344
Lys Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr	
435 440 445	
tgc aac ctg aag cgg tgc tca gag aca gga ggg agt gtt gtg gaa ttg	1392
Cys Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly Ser Val Val Glu Leu	
450 455 460	
ccc aca gtt tcc cag gaa cca agt ggg ccg agc gac tct gag aca gac	1440
Pro Thr Val Ser Gln Glu Pro Ser Gly Pro Ser Asp Ser Glu Thr Asp	
465 470 475 480	
tgc atg tat ggg aat ggc aaa gac tat cgg ggc aaa acg gcc gtc act	1488
Cys Met Tyr Gly Asn Gly Lys Asp Tyr Arg Gly Lys Thr Ala Val Thr	
485 490 495	
gca gct ggc acc ccc tgc cag gga tgg gct gcc cag gag ccc cac agg	1536
Ala Ala Gly Thr Pro Cys Gln Gly Trp Ala Ala Gln Glu Pro His Arg	
500 505 510	
cac agc atc ttc acc cca cag aca aac cca cgg gca gat ctg gaa aag	1584
His Ser Ile Phe Thr Pro Gln Thr Asn Pro Arg Ala Asp Leu Glu Lys	
515 520 525	
aac tac tgc cga aac cca gat ggg gat gtg aat ggt cct tgg tgc tat	1632
Asn Tyr Cys Arg Asn Pro Asp Gly Asp Val Asn Gly Pro Trp Cys Tyr	
530 535 540	
aca aca aac ccc aga aaa ctt tat gac tat tgt gac atc ccc ctg tgt	1680
Thr Thr Asn Pro Arg Lys Leu Tyr Asp Tyr Cys Asp Ile Pro Leu Cys	
545 550 555 560	

gca tca gca tca tcc ttt gag tgc ggg aaa cct cag gtg gaa ccg aag	1728
Ala Ser Ala Ser Ser Phe Glu Cys Gly Lys Pro Gln Val Glu Pro Lys	
565 570 575	
aaa tgc cct ggg agg gtg gtg ggt ggc tgc gtg gcc aac cct cac tcc	1776
Lys Cys Pro Gly Arg Val Val Gly Gly Cys Val Ala Asn Pro His Ser	
580 585 590	
tgg ccc tgg caa atc agc ctt aga aca aga ttt acc gga cag cac ttc	1824
Trp Pro Trp Gln Ile Ser Leu Arg Thr Arg Phe Thr Gly Gln His Phe	
595 600 605	
tgt ggc ggt act tta ata gcc cca gag tgg gtt ctg act gct gcc cac	1872
Cys Gly Gly Thr Leu Ile Ala Pro Glu Trp Val Leu Thr Ala Ala His	
610 615 620	
tgt ttg gag aaa tct tca aga cct gaa ttc tac aag gtt atc ctg ggt	1920
Cys Leu Glu Lys Ser Ser Arg Pro Glu Phe Tyr Lys Val Ile Leu Gly	
625 630 635 640	
gcg cac gaa gaa tat atc cgt ggg ttg gat gtt cag gaa ata tca gta	1968
Ala His Glu Glu Tyr Ile Arg Gly Leu Asp Val Gln Glu Ile Ser Val	
645 650 655	
gcc aaa ctg atc ttg gag ccc aac aac cgt gac att gcc ctg ctg aaa	2016
Ala Lys Leu Ile Leu Glu Pro Asn Asn Arg Asp Ile Ala Leu Leu Lys	
660 665 670	
cta agc cgc cca gcc acc atc acg gat aaa gtc att cca gct tgt ctg	2064
Leu Ser Arg Pro Ala Thr Ile Thr Asp Lys Val Ile Pro Ala Cys Leu	
675 680 685	
cca tct cca aat tac atg gtt gct gac cgg aca ata tgt tac atc acc	2112
Pro Ser Pro Asn Tyr Met Val Ala Asp Arg Thr Ile Cys Tyr Ile Thr	
690 695 700	
ggc tgg gga gag act caa ggg act ttc ggt gcc ggt cgt ctc aag gag	2160
Gly Trp Gly Glu Thr Gln Gly Thr Phe Gly Ala Gly Arg Leu Lys Glu	
705 710 715 720	
gct cag ctg cct gtg att gag aac aag gtg tgc aac cgc gtc gag tat	2208
Ala Gln Leu Pro Val Ile Glu Asn Lys Val Cys Asn Arg Val Glu Tyr	
725 730 735	
ctg aac aac aga gtc aaa tcc acg gag ctc tgt gcc ggg caa ctg gct	2256
Leu Asn Asn Arg Val Lys Ser Thr Glu Leu Cys Ala Gly Gln Leu Ala	
740 745 750	
ggc ggc gtc gac agc tgc caa ggc gac agt gga gga cct ctg gtt tgc	2304
Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys	
755 760 765	
ttc gag aag gac aag tac att tta caa gga gtc act tct tgg ggt ctt	2352
Phe Glu Lys Asp Lys Tyr Ile Leu Gln Gly Val Thr Ser Trp Gly Leu	
770 775 780	

ggc tgt gct cgc ccc aat aag cct ggt gtc tac gtt cgt gtc tca cgg 2400
 Gly Cys Ala Arg Pro Asn Lys Pro Gly Val Tyr Val Arg Val Ser Arg
 785 790 795 800

ttt gtt gat tgg att gaa agg gag atg agg aat aac tgactaggtg 2446
 Phe Val Asp Trp Ile Glu Arg Glu Met Arg Asn Asn
 805 810

gaaggccgag caaaacctct gcttactaaa gcttactgaa tatggggaga gggcttaggg 2506
 tgtttgga aactgacagt aatcaaaactg ggacactaca ctgaaccaca gcttctgtgc 2566
 gccctcagc ccctccctt tttttgtatt attgtgggta aaattttcct gtctgtggac 2626
 ttctggattt tgtgacaata gaccatcact gctgtgacct ttgttgaaaa taaactcgat 2686
 acttactttg 2696

<210> 4
 <211> 812
 <212> PRT
 <213> Mus musculus

<400> 4
 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro
 1 5 10 15
 Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser
 20 25 30
 Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp
 35 40 45
 Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe
 50 55 60
 Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Ser
 65 70 75 80
 Lys Thr Ser Ser Ile Ile Arg Met Arg Asp Val Ile Leu Phe Glu Lys
 85 90 95
 Arg Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg
 100 105 110
 Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly
 115 120 125
 Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn
 130 135 140
 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln
 145 150 155 160
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys
 165 170 175
 Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys
 180 185 190
 Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala
 195 200 205
 Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe
 210 215 220
 Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu
 225 230 235 240
 Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr
 245 250 255
 Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr
 260 265 270

Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser
 275 280 285
 Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro
 290 295 300
 His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu
 305 310 315 320
 Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr
 325 330 335
 Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys
 340 345 350
 Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu
 355 360 365
 Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser
 370 375 380
 Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser
 385 390 395 400
 Trp Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe
 405 410 415
 Pro Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp
 420 425 430
 Lys Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr
 435 440 445
 Cys Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly Ser Val Val Glu Leu
 450 455 460
 Pro Thr Val Ser Gln Glu Pro Ser Gly Pro Ser Asp Ser Glu Thr Asp
 465 470 475 480
 Cys Met Tyr Gly Asn Gly Lys Asp Tyr Arg Gly Lys Thr Ala Val Thr
 485 490 495
 Ala Ala Gly Thr Pro Cys Gln Gly Trp Ala Ala Gln Glu Pro His Arg
 500 505 510
 His Ser Ile Phe Thr Pro Gln Thr Asn Pro Arg Ala Asp Leu Glu Lys
 515 520 525
 Asn Tyr Cys Arg Asn Pro Asp Gly Asp Val Asn Gly Pro Trp Cys Tyr
 530 535 540
 Thr Thr Asn Pro Arg Lys Leu Tyr Asp Tyr Cys Asp Ile Pro Leu Cys
 545 550 555 560
 Ala Ser Ala Ser Ser Phe Glu Cys Gly Lys Pro Gln Val Glu Pro Lys
 565 570 575
 Lys Cys Pro Gly Arg Val Val Gly Gly Cys Val Ala Asn Pro His Ser
 580 585 590
 Trp Pro Trp Gln Ile Ser Leu Arg Thr Arg Phe Thr Gly Gln His Phe
 595 600 605
 Cys Gly Gly Thr Leu Ile Ala Pro Glu Trp Val Leu Thr Ala Ala His
 610 615 620
 Cys Leu Glu Lys Ser Ser Arg Pro Glu Phe Tyr Lys Val Ile Leu Gly
 625 630 635 640
 Ala His Glu Glu Tyr Ile Arg Gly Leu Asp Val Gln Glu Ile Ser Val
 645 650 655
 Ala Lys Leu Ile Leu Glu Pro Asn Asn Arg Asp Ile Ala Leu Leu Lys
 660 665 670
 Leu Ser Arg Pro Ala Thr Ile Thr Asp Lys Val Ile Pro Ala Cys Leu
 675 680 685
 Pro Ser Pro Asn Tyr Met Val Ala Asp Arg Thr Ile Cys Tyr Ile Thr
 690 695 700
 Gly Trp Gly Glu Thr Gln Gly Thr Phe Gly Ala Gly Arg Leu Lys Glu


```

705              710              715              720
Ala Gln Leu Pro Val Ile Glu Asn Lys Val Cys Asn Arg Val Glu Tyr
              725              730              735
Leu Asn Asn Arg Val Lys Ser Thr Glu Leu Cys Ala Gly Gln Leu Ala
              740              745              750
Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys
              755              760              765
Phe Glu Lys Asp Lys Tyr Ile Leu Gln Gly Val Thr Ser Trp Gly Leu
              770              775              780
Gly Cys Ala Arg Pro Asn Lys Pro Gly Val Tyr Val Arg Val Ser Arg
785              790              795              800
Phe Val Asp Trp Ile Glu Arg Glu Met Arg Asn Asn
              805              810

```

```

<210> 5
<211> 1083
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (1)...(1083)

```

```

<400> 5
tat ctc tca gag tgc aag act ggg aat gga aag aac tac aga ggg acg      48
Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg Gly Thr
  1              5              10              15

atg tcc aaa aca aaa aat ggc atc acc tgt caa aaa tgg agt tcc act      96
Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser Ser Thr
      20              25              30

tct ccc cac aga cct aga ttc tca cct gct aca cac ccc tca gag gga      144
Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser Glu Gly
      35              40              45

ctg gag gag aac tac tgc agg aat cca gac aac gat ccg cag ggg ccc      192
Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln Gly Pro
      50              55              60

tgg tgc tat act act gat cca gaa aag aga tat gac tac tgc gac att      240
Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys Asp Ile
      65              70              75              80

ctt gag tgt gaa gag gaa tgt atg cat tgc agt gga gaa aac tat gac      288
Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn Tyr Asp
      85              90              95

ggc aaa att tcc aag acc atg tct gga ctg gaa tgc cag gcc tgg gac      336
Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala Trp Asp
      100              105              110

tct cag agc cca cac gct cat gga tac att cct tcc aaa ttt cca aac      384
Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe Pro Asn
      115              120              125

```

aag aac ctg aag aag aat tac tgt cgt aac ccc gat agg gag ctg cgg Lys Asn Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu Leu Arg 130 135 140	432
cct tgg tgt ttc acc acc gac ccc aac aag cgc tgg gaa ctt tgc gac Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu Cys Asp 145 150 155 160	480
atc ccc cgc tgc aca aca cct cca cca tct tct ggt ccc acc tac cag Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Ser Gly Pro Thr Tyr Gln 165 170 175	528
tgt ctg aag gga aca ggt gaa aac tat cgc ggg aat gtg gct gtt acc Cys Leu Lys Gly Thr Glu Asn Tyr Arg Gly Asn Val Ala Val Thr 180 185 190	576
gtt tcc ggg cac acc tgt cag cac tgg agt gca cag acc cct cac aca Val Ser Gly His Thr Cys Gln His Trp Ser Ala Gln Thr Pro His Thr 195 200 205	624
cat aac agg aca cca gaa aac ttc ccc tgc aaa aat ttg gat gaa aac His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Asp Glu Asn 210 215 220	672
tac tgc cgc aat cct gac gga aaa agg gcc cca tgg tgc cat aca acc Tyr Cys Arg Asn Pro Asp Gly Lys Arg Ala Pro Trp Cys His Thr Thr 225 230 235 240	720
aac agc caa gtg cgg tgg gag tac tgt aag ata ccg tcc tgt gac tcc Asn Ser Gln Val Arg Trp Glu Tyr Cys Lys Ile Pro Ser Cys Asp Ser 245 250 255	768
tcc cca gta tcc acg gaa caa ttg gct ccc aca gca cca cct gag cta Ser Pro Val Ser Thr Glu Gln Leu Ala Pro Thr Ala Pro Pro Glu Leu 260 265 270	816
acc cct gtg gtc cag gac tgc tac cat ggt gat gga cag agc tac cga Thr Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Gln Ser Tyr Arg 275 280 285	864
ggc aca tcc tcc acc acc acc aca gga aag aag tgt cag tct tgg tca Gly Thr Ser Ser Thr Thr Thr Thr Gly Lys Lys Cys Gln Ser Trp Ser 290 295 300	912
tct atg aca cca cac cgg cac cag aag acc cca gaa aac tac cca aat Ser Met Thr Pro His Arg His Gln Lys Thr Pro Glu Asn Tyr Pro Asn 305 310 315 320	960
gct ggc ctg aca atg aac tac tgc agg aat cca gat gcc gat aaa ggc Ala Gly Leu Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp Lys Gly 325 330 335	1008
ccc tgg tgt ttt acc aca gac ccc agc gtc agg tgg gag tac tgc aac Pro Trp Cys Phe Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr Cys Asn	1056

340 345 350
 ctg aaa aaa tgc tca gga aca gaa gcg 1083
 Leu Lys Lys Cys Ser Gly Thr Glu Ala
 355 360

<210> 6
 <211> 361
 <212> PRT
 <213> Homo sapiens

<400> 6
 Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg Gly Thr
 1 5 10 15
 Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser Ser Thr
 20 25 30
 Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser Glu Gly
 35 40 45
 Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln Gly Pro
 50 55 60
 Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys Asp Ile
 65 70 75 80
 Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn Tyr Asp
 85 90 95
 Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala Trp Asp
 100 105 110
 Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe Pro Asn
 115 120 125
 Lys Asn Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu Leu Arg
 130 135 140
 Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu Cys Asp
 145 150 155 160
 Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Gly Pro Thr Tyr Gln
 165 170 175
 Cys Leu Lys Gly Thr Gly Glu Asn Tyr Arg Gly Asn Val Ala Val Thr
 180 185 190
 Val Ser Gly His Thr Cys Gln His Trp Ser Ala Gln Thr Pro His Thr
 195 200 205
 His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Asp Glu Asn
 210 215 220
 Tyr Cys Arg Asn Pro Asp Gly Lys Arg Ala Pro Trp Cys His Thr Thr
 225 230 235 240
 Asn Ser Gln Val Arg Trp Glu Tyr Cys Lys Ile Pro Ser Cys Asp Ser
 245 250 255
 Ser Pro Val Ser Thr Glu Gln Leu Ala Pro Thr Ala Pro Pro Glu Leu
 260 265 270
 Thr Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Gln Ser Tyr Arg
 275 280 285
 Gly Thr Ser Ser Thr Thr Thr Thr Gly Lys Lys Cys Gln Ser Trp Ser
 290 295 300
 Ser Met Thr Pro His Arg His Gln Lys Thr Pro Glu Asn Tyr Pro Asn
 305 310 315 320
 Ala Gly Leu Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp Lys Gly
 325 330 335

Pro Trp Cys Phe Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr Cys Asn
 340 345 350
 Leu Lys Lys Cys Ser Gly Thr Glu Ala
 355 360

<210> 7
 <211> 1086
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)...(1086)

<400> 7
 gtg tat ctg tca gaa tgt aag acc ggc atc ggc aac ggc tac aga gga 48
 Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg Gly
 1 5 10 15
 acc atg tcc agg aca aag agt ggt gtt gcc tgt caa aag tgg ggt gcc 96
 Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly Ala
 20 25 30
 acg ttc ccc cac gta ccc aac tac tct ccc agt aca cat ccc aat gag 144
 Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn Glu
 35 40 45
 gga cta gaa gag aac tac tgt agg aac cca gac aat gat gaa caa ggg 192
 Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln Gly
 50 55 60
 cct tgg tgc tac act aca gat ccg gac aag aga tat gac tac tgc aac 240
 Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys Asn
 65 70 75 80
 att cct gaa tgt gaa gag gaa tgc atg tac tgc agt gga gaa aag tat 288
 Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys Tyr
 85 90 95
 gag ggc aaa atc tcc aag acc atg tct gga ctt gac tgc cag gcc tgg 336
 Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala Trp
 100 105 110
 gat tct cag agc cca cat gct cat gga tac atc cct gcc aaa ttt cca 384
 Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe Pro
 115 120 125
 agc aag aac ctg aag atg aat tat tgc cac aac cct gac ggg gag cca 432
 Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu Pro
 130 135 140
 agg ccc tgg tgc ttc aca aca gac ccc acc aaa cgc tgg gaa tac tgt 480
 Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr Cys
 145 150 155 160

gac atc ccc cgc tgc aca aca ccc ccg ccc cca ccc agc cca acc tac Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Pro Ser Pro Thr Tyr 165 170 175	528
caa tgt ctg aaa gga aga ggt gaa aat tac cga ggg acc gtg tct gtc Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser Val 180 185 190	576
acc gtg tct ggg aaa acc tgt cag cgc tgg agt gag caa acc cct cat Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro His 195 200 205	624
agg cac aac agg aca cca gaa aat ttc ccc tgc aaa aat ctg gaa gag Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu Glu 210 215 220	672
aac tac tgc cgg aac cca gat gga gaa act gct ccc tgg tgc tat acc Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr Thr 225 230 235 240	720
act gac agc cag ctg agg tgg gag tac tgt gag att cca tcc tgc gag Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys Glu 245 250 255	768
tcc tca gca tca cca gac cag tca gat tcc tca gtt cca cca gag gag Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu Glu 260 265 270	816
caa aca cct gtg gtc cag gaa tgc tac cag agc gat ggg cag agc tat Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser Tyr 275 280 285	864
cgg ggt aca tcg tcc act acc atc aca ggg aag aag tgc cag tcc tgg Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser Trp 290 295 300	912
gca gct atg ttt cca cac agg cat tcg aag acc cca gag aac ttc cca Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe Pro 305 310 315 320	960
gat gct ggc ttg gag atg aac tac tgc agg aac ccg gat ggt gac aag Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp Lys 325 330 335	1008
ggc cct tgg tgc tac acc act gac ccg agc gtc agg tgg gaa tac tgc Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr Cys 340 345 350	1056
aac ctg aag cgg tgc tca gag aca gga ggg Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly 355 360	1086

<210> 8

<211> 362

<212> PRT

<213> Mus musculus

<400> 8

Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg Gly
 1 5 10 15
 Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly Ala
 20 25 30
 Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn Glu
 35 40 45
 Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln Gly
 50 55 60
 Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys Asn
 65 70 75 80
 Ile Pro Glu Cys Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys Tyr
 85 90 95
 Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala Trp
 100 105 110
 Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe Pro
 115 120 125
 Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu Pro
 130 135 140
 Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr Cys
 145 150 155 160
 Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr Tyr
 165 170 175
 Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser Val
 180 185 190
 Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro His
 195 200 205
 Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu Glu
 210 215 220
 Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr Thr
 225 230 235 240
 Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys Glu
 245 250 255
 Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu Glu
 260 265 270
 Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser Tyr
 275 280 285
 Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser Trp
 290 295 300
 Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe Pro
 305 310 315 320
 Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp Lys
 325 330 335
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr Cys
 340 345 350
 Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly
 355 360

<210> 9

<211> 552

<212> DNA

<213> Mus musculus

<220>
 <221> CDS
 <222> (1) ... (552)

<400> 9
 cat act cat cag gac ttt cag cca gtg ctc cac ctg gtg gca ctg aac 48
 His Thr His Gln Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn
 1 5 10 15
 acc ccc ctg tct gga ggc atg cgt ggt atc cgt gga gca gat ttc cag 96
 Thr Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln
 20 25 30
 tgc ttc cag caa gcc cga gcc gtg ggg ctg tcg ggc acc ttc cgg gct 144
 Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ser Gly Thr Phe Arg Ala
 35 40 45
 ttc ctg tcc tct agg ctg cag gat ctc tat agc atc gtg cgc cgt gct 192
 Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala
 50 55 60
 gac cgg ggg tct gtg ccc atc gtc aac ctg aag gac gag gtg cta tct 240
 Asp Arg Gly Ser Val Pro Ile Val Asn Leu Lys Asp Glu Val Leu Ser
 65 70 75 80
 ccc agc tgg gac tcc ctg ttt tct ggc tcc cag ggt caa gtg caa ccc 288
 Pro Ser Trp Asp Ser Leu Phe Ser Gly Ser Gln Gly Gln Val Gln Pro
 85 90 95
 ggg gcc cgc atc ttt tct ttt gac ggc aga gat gtc ctg aga cac cca 336
 Gly Ala Arg Ile Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His Pro
 100 105 110
 gcc tgg ccg cag aag agc gta tgg cac ggc tcg gac ccc agt ggg cgg 384
 Ala Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Ser Gly Arg
 115 120 125
 agg ctg atg gag agt tac tgt gag aca tgg cga act gaa act act ggg 432
 Arg Leu Met Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr Gly
 130 135 140
 gct aca ggt cag gcc tcc tcc ctg ctg tca ggc agg ctc ctg gaa cag 480
 Ala Thr Gly Gln Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu Gln
 145 150 155 160
 aaa gct gcg agc tgc cac aac agc tac atc gtc ctg tgc att gag aat 528
 Lys Ala Ala Ser Cys His Asn Ser Tyr Ile Val Leu Cys Ile Glu Asn
 165 170 175
 agc ttc atg acc tct ttc tcc aaa 552
 Ser Phe Met Thr Ser Phe Ser Lys
 180

<210> 10

<211> 184
 <212> PRT
 <213> Mus musculus

<400> 10
 His Thr His Gln Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn
 1 5 10 15
 Thr Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln
 20 25 30
 Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ser Gly Thr Phe Arg Ala
 35 40 45
 Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala
 50 55 60
 Asp Arg Gly Ser Val Pro Ile Val Asn Leu Lys Asp Glu Val Leu Ser
 65 70 75 80
 Pro Ser Trp Asp Ser Leu Phe Ser Gly Ser Gln Gly Gln Val Gln Pro
 85 90 95
 Gly Ala Arg Ile Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His Pro
 100 105 110
 Ala Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Ser Gly Arg
 115 120 125
 Arg Leu Met Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr Gly
 130 135 140
 Ala Thr Gly Gln Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu Gln
 145 150 155 160
 Lys Ala Ala Ser Cys His Asn Ser Tyr Ile Val Leu Cys Ile Glu Asn
 165 170 175
 Ser Phe Met Thr Ser Phe Ser Lys
 180

<210> 11
 <211> 1414
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)...(1414)

<400> 11
 atg gac cat aag gaa gta atc ctt ctg ttt ctc ttg ctt ctg aaa cca 48
 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro
 1 5 10 15
 gga caa ggg gac tcg ctg gat ggc tac ata agc aca caa ggg gct tca 96
 Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser
 20 25 30
 ctg ttc agt ctc acc aag aag cag ctc gca gca gga ggt gtc tcg gac 144
 Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp
 35 40 45
 tgt ttg gcc aaa tgt gaa ggg gaa aca gac ttt gtc tgc agg tca ttc 192
 Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe
 50 55 60

cag tac cac agc aaa gag cag caa tgc gtg atc atg gcg gag aac agc	240
Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Ser	
65 70 75 80	
aag act tcc tcc atc atc cgg atg aga gac gtc atc tta ttc gaa aag	288
Lys Thr Ser Ser Ile Ile Arg Met Arg Asp Val Ile Leu Phe Glu Lys	
85 90 95	
aga gtg tat ctg tca gaa tgt aag acc ggc atc ggc aac ggc tac aga	336
Arg Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg	
100 105 110	
gga acc atg tcc agg aca aag agt ggt gtt gcc tgt caa aag tgg ggt	384
Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly	
115 120 125	
gcc acg ttc ccc cac gta ccc aac tac tct ccc agt aca cat ccc aat	432
Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn	
130 135 140	
gag gga cta gaa gag aac tac tgt agg aac cca gac aat gat gaa caa	480
Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln	
145 150 155 160	
ggg cct tgg tgc tac act aca gat ccg gac aag aga tat gac tac tgc	528
Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys	
165 170 175	
aac att cct gaa tgt gaa gag gaa tgc atg tac tgc agt gga gaa aag	576
Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys	
180 185 190	
tat gag ggc aaa atc tcc aag acc atg tct gga ctt gac tgc cag gcc	624
Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala	
195 200 205	
tgg gat tct cag agc cca cat gct cat gga tac atc cct gcc aaa ttt	672
Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe	
210 215 220	
cca agc aag aac ctg aag atg aat tat tgc cac aac cct gac ggg gag	720
Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu	
225 230 235 240	
cca agg ccc tgg tgc ttc aca aca gac ccc acc aaa cgc tgg gaa tac	768
Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr	
245 250 255	
tgt gac atc ccc cgc tgc aca aca ccc ccg ccc cca ccc agc cca acc	816
Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr	
260 265 270	
tac caa tgt ctg aaa gga aga ggt gaa aat tac cga ggg acc gtg tct	864
Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser	

275	280	285	
gtc acc gtg tct ggg aaa acc tgt cag cgc tgg agt gag caa acc cct			912
Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro			
290	295	300	
cat agg cac aac agg aca cca gaa aat ttc ccc tgc aaa aat ctg gaa			960
His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu			
305	310	315	320
gag aac tac tgc cgg aac cca gat gga gaa act gct ccc tgg tgc tat			1008
Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr			
	325	330	335
acc act gac agc cag ctg agg tgg gag tac tgt gag att cca tcc tgc			1056
Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys			
	340	345	350
gag tcc tca gca tca cca gac cag tca gat tcc tca gtt cca cca gag			1104
Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu			
	355	360	365
gag caa aca cct gtg gtc cag gaa tgc tac cag agc gat ggg cag agc			1152
Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser			
	370	375	380
tat cgg ggt aca tcg tcc act acc atc aca ggg aag aag tgc cag tcc			1200
Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser			
	385	390	395
400			
tgg gca gct atg ttt cca cac agg cat tcg aag acc cca gag aac ttc			1248
Trp Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe			
	405	410	415
cca gat gct ggc ttg gag atg aac tac tgc agg aac ccg gat ggt gac			1296
Pro Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp			
	420	425	430
aag ggc cct tgg tgc tac acc act gac ccg agc gtc agg tgg gaa tac			1344
Lys Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr			
	435	440	445
tgc aac ctg aag cgg tgc tca gag aca gga ggg aat tca gac tac aag			1392
Cys Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly Asn Ser Asp Tyr Lys			
	450	455	460
gac gac gat gac aag taa taa c			1414
Asp Asp Asp Asp Lys * *			
465			

<210> 12
 <211> 469
 <212> PRT
 <213> Mus musculus

<400> 12
 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro
 1 5 10 15
 Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser
 20 25 30
 Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp
 35 40 45
 Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe
 50 55 60
 Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Ser
 65 70 75 80
 Lys Thr Ser Ser Ile Ile Arg Met Arg Asp Val Ile Leu Phe Glu Lys
 85 90 95
 Arg Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg
 100 105 110
 Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly
 115 120 125
 Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn
 130 135 140
 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln
 145 150 155 160
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys
 165 170 175
 Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys
 180 185 190
 Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala
 195 200 205
 Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe
 210 215 220
 Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu
 225 230 235 240
 Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr
 245 250 255
 Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr
 260 265 270
 Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser
 275 280 285
 Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro
 290 295 300
 His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu
 305 310 315 320
 Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr
 325 330 335
 Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys
 340 345 350
 Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu
 355 360 365
 Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser
 370 375 380
 Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser
 385 390 395 400
 Trp Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe
 405 410 415
 Pro Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp
 420 425 430

Lys Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr
 435 440 445
 Cys Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly Asn Ser Asp Tyr Lys
 450 455 460
 Asp Asp Asp Asp Lys
 465

<210> 13
 <211> 661
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)...(661)

<400> 13
 atg gac cat aag gaa gta atc ctt ctg ttt ctc ttg ctt ctg aaa cca 48
 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro
 1 5 10 15
 gga caa ggg gac tcg cta gat ctt gac tac aag gac gac gat gac aag 96
 Gly Gln Gly Asp Ser Leu Asp Leu Asp Tyr Lys Asp Asp Asp Lys
 20 25 30
 ctt gct cat act cat cag gac ttt cag cca gtg ctc cac ctg gtg gca 144
 Leu Ala His Thr His Gln Asp Phe Gln Pro Val Leu His Leu Val Ala
 35 40 45
 ctg aac acc ccc ctg tct gga ggc atg cgt ggt atc cgt gga gca gat 192
 Leu Asn Thr Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp
 50 55 60
 ttc cag tgc ttc cag caa gcc cga gcc gtg ggg ctg tcg ggc acc ttc 240
 Phe Gln Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ser Gly Thr Phe
 65 70 75 80
 cgg gct ttc ctg tcc tct agg ctg cag gat ctc tat agc atc gtg cgc 288
 Arg Ala Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg
 85 90 95
 cgt gct gac cgg ggg tct gtg ccc atc gtc aac ctg aag gac gag gtg 336
 Arg Ala Asp Arg Gly Ser Val Pro Ile Val Asn Leu Lys Asp Glu Val
 100 105 110
 cta tct ccc agc tgg gac tcc ctg ttt tct ggc tcc cag ggt caa gtg 384
 Leu Ser Pro Ser Trp Asp Ser Leu Phe Ser Gly Ser Gln Gly Gln Val
 115 120 125
 caa ccc ggg gcc cgc atc ttt tct ttt gac ggc aga gat gtc ctg aga 432
 Gln Pro Gly Ala Arg Ile Phe Ser Phe Asp Gly Arg Asp Val Leu Arg
 130 135 140
 cac cca gcc tgg ccg cag aag agc gta tgg cac ggc tcg gac ccc agt 480
 His Pro Ala Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Ser

145	150	155	160	
ggg cgg agg ctg atg gag agt tac tgt gag aca tgg cga act gaa act				528
Gly Arg Arg Leu Met Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr				
165		170	175	
act ggg gct aca ggt cag gcc tcc tcc ctg ctg tca ggc agg ctc ctg				576
Thr Gly Ala Thr Gly Gln Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu				
180	185	190		
gaa cag aaa gct gcg agc tgc cac aac agc tac atc gtc ctg tgc att				624
Glu Gln Lys Ala Ala Ser Cys His Asn Ser Tyr Ile Val Leu Cys Ile				
195	200	205		
gag aat agc ttc atg acc tct ttc tcc aaa taa taa c				661
Glu Asn Ser Phe Met Thr Ser Phe Ser Lys * *				
210	215			

<210> 14
 <211> 218
 <212> PRT
 <213> Mus musculus

<400> 14

Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Leu Lys Pro				
1	5	10	15	
Gly Gln Gly Asp Ser Leu Asp Leu Asp Tyr Lys Asp Asp Asp Asp Lys				
20	25	30		
Leu Ala His Thr His Gln Asp Phe Gln Pro Val Leu His Leu Val Ala				
35	40	45		
Leu Asn Thr Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp				
50	55	60		
Phe Gln Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ser Gly Thr Phe				
65	70	75	80	
Arg Ala Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg				
85	90	95		
Arg Ala Asp Arg Gly Ser Val Pro Ile Val Asn Leu Lys Asp Glu Val				
100	105	110		
Leu Ser Pro Ser Trp Asp Ser Leu Phe Ser Gly Ser Gln Gly Gln Val				
115	120	125		
Gln Pro Gly Ala Arg Ile Phe Ser Phe Asp Gly Arg Asp Val Leu Arg				
130	135	140		
His Pro Ala Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Ser				
145	150	155	160	
Gly Arg Arg Leu Met Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr				
165	170	175		
Thr Gly Ala Thr Gly Gln Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu				
180	185	190		
Glu Gln Lys Ala Ala Ser Cys His Asn Ser Tyr Ile Val Leu Cys Ile				
195	200	205		
Glu Asn Ser Phe Met Thr Ser Phe Ser Lys				
210	215			

<210> 15

<211> 1747
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)...(1747)

<400> 15
 atg gac cat aag gaa gta atc ctt ctg ttt ctc ttg ctt ctg aaa cca 48
 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro
 1 5 10 15

gga caa ggg gac tcg ctg gat ggc tac ata agc aca caa ggg gct tca 96
 Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser
 20 25 30

ctg ttc agt ctc acc aag aag cag ctc gca gca gga ggt gtc tcg gac 144
 Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp
 35 40 45

tgt ttg gcc aaa tgt gaa ggg gaa aca gac ttt gtc tgc agg tca ttc 192
 Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe
 50 55 60

cag tac cac agc aaa gag cag caa tgc gtg atc atg gcg gag aac agc 240
 Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Ser
 65 70 75 80

aag act tcc tcc atc atc cgg atg aga gac gtc atc tta ttc gaa aag 288
 Lys Thr Ser Ser Ile Ile Arg Met Arg Asp Val Ile Leu Phe Glu Lys
 85 90 95

aga gtg tat ctg tca gaa tgt aag acc ggc atc ggc aac ggc tac aga 336
 Arg Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg
 100 105 110

gga acc atg tcc agg aca aag agt ggt gtt gcc tgt caa aag tgg ggt 384
 Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly
 115 120 125

gcc acg ttc ccc cac gta ccc aac tac tct ccc agt aca cat ccc aat 432
 Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn
 130 135 140

gag gga cta gaa gag aac tac tgt agg aac cca gac aat gat gaa caa 480
 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln
 145 150 155 160

ggg cct tgg tgc tac act aca gat ccg gac aag aga tat gac tac tgc 528
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Lys Arg Tyr Asp Tyr Cys
 165 170 175

aac att cct gaa tgt gaa gag gaa tgc atg tac tgc agt gga gaa aag 576
 Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys

180	185	190	
tat gag ggc aaa atc tcc aag acc atg tct gga ctt gac tgc cag gcc			624
Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala			
195	200	205	
tgg gat tct cag agc cca cat gct cat gga tac atc cct gcc aaa ttt			672
Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe			
210	215	220	
cca agc aag aac ctg aag atg aat tat tgc cac aac cct gac ggg gag			720
Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu			
225	230	235	240
cca agg ccc tgg tgc ttc aca aca gac ccc acc aaa cgc tgg gaa tac			768
Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr			
245	250	255	
tgt gac atc ccc cgc tgc aca aca ccc ccg ccc cca ccc agc cca acc			816
Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Ser Pro Thr			
260	265	270	
tac caa tgt ctg aaa gga aga ggt gaa aat tac cga ggg acc gtg tct			864
Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser			
275	280	285	
gtc acc gtg tct ggg aaa acc tgt cag cgc tgg agt gag caa acc cct			912
Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro			
290	295	300	
cat agg cac aac agg aca cca gaa aat ttc ccc tgc aaa aat ctg gaa			960
His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu			
305	310	315	320
gag aac tac tgc cgg aac cca gat gga gaa act gct ccc tgg tgc tat			1008
Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr			
325	330	335	
acc act gac agc cag ctg agg tgg gag tac tgt gag att cca tcc tgc			1056
Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys			
340	345	350	
gag tcc tca gca tca cca gac cag tca gat tcc tca gtt cca cca gag			1104
Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu			
355	360	365	
gag caa aca cct gtg gga ggg aat tgc ggc ggt gga tca ggt ggc gga			1152
Glu Gln Thr Pro Val Gly Gly Asn Cys Gly Gly Gly Ser Gly Gly Gly			
370	375	380	
gat ctt gac tac aag gac gac gat gac aag ctt gct cat act cat cag			1200
Asp Leu Asp Tyr Lys Asp Asp Asp Asp Lys Leu Ala His Thr His Gln			
385	390	395	400
gac ttt cag cca gtg ctc cac ctg gtg gca ctg aac acc ccc ctg tct			1248

Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn Thr Pro Leu Ser
 405 410 415
 gga ggc atg cgt ggt atc cgt gga gca gat ttc cag tgc ttc cag caa 1296
 Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln Cys Phe Gln Gln
 420 425 430
 gcc cga gcc gtg ggg ctg tcg ggc acc ttc cgg gct ttc ctg tcc tct 1344
 Ala Arg Ala Val Gly Leu Ser Gly Thr Phe Arg Ala Phe Leu Ser Ser
 435 440 445
 agg ctg cag gat ctc tat agc atc gtg cgc cgt gct gac cgg ggg tct 1392
 Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala Asp Arg Gly Ser
 450 455 460
 gtg ccc atc gtc aac ctg aag gac gag gtg cta tct ccc agc tgg gac 1440
 Val Pro Ile Val Asn Leu Lys Asp Glu Val Leu Ser Pro Ser Trp Asp
 465 470 475 480
 tcc ctg ttt tct ggc tcc cag ggt caa gtg caa ccc ggg gcc cgc atc 1488
 Ser Leu Phe Ser Gly Ser Gln Gly Gln Val Gln Pro Gly Ala Arg Ile
 485 490 495
 ttt tct ttt gac ggc aga gat gtc ctg aga cac cca gcc tgg ccg cag 1536
 Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His Pro Ala Trp Pro Gln
 500 505 510
 aag agc gta tgg cac ggc tcg gac ccc agt ggg cgg agg ctg atg gag 1584
 Lys Ser Val Trp His Gly Ser Asp Pro Ser Gly Arg Arg Leu Met Glu
 515 520 525
 agt tac tgt gag aca tgg cga act gaa act act ggg gct aca ggt cag 1632
 Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr Gly Ala Thr Gly Gln
 530 535 540
 gcc tcc tcc ctg ctg tca ggc agg ctc ctg gaa cag aaa gct gcg agc 1680
 Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu Gln Lys Ala Ala Ser
 545 550 555 560
 tgc cac aac agc tac atc gtc ctg tgc att gag aat agc ttc atg acc 1728
 Cys His Asn Ser Tyr Ile Val Leu Cys Ile Glu Asn Ser Phe Met Thr
 565 570 575
 tct ttc tcc aaa taa taa c 1747
 Ser Phe Ser Lys * *
 580

<210> 16
 <211> 580
 <212> PRT
 <213> Mus musculus

<400> 16
 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Lys Pro

1	5	10	15
Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser			
20	25	30	
Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp			
35	40	45	
Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe			
50	55	60	
Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Ser			
65	70	75	80
Lys Thr Ser Ser Ile Ile Arg Met Arg Asp Val Ile Leu Phe Glu Lys			
85	90	95	
Arg Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg			
100	105	110	
Gly Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly			
115	120	125	
Ala Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn			
130	135	140	
Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln			
145	150	155	160
Gly Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys			
165	170	175	
Asn Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys			
180	185	190	
Tyr Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala			
195	200	205	
Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe			
210	215	220	
Pro Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu			
225	230	235	240
Pro Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr			
245	250	255	
Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Pro Ser Pro Thr			
260	265	270	
Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser			
275	280	285	
Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro			
290	295	300	
His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu			
305	310	315	320
Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr			
325	330	335	
Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys			
340	345	350	
Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu			
355	360	365	
Glu Gln Thr Pro Val Gly Gly Asn Cys Gly Gly Gly Ser Gly Gly Gly			
370	375	380	
Asp Leu Asp Tyr Lys Asp Asp Asp Asp Lys Leu Ala His Thr His Gln			
385	390	395	400
Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn Thr Pro Leu Ser			
405	410	415	
Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln Cys Phe Gln Gln			
420	425	430	
Ala Arg Ala Val Gly Leu Ser Gly Thr Phe Arg Ala Phe Leu Ser Ser			
435	440	445	

Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala Asp Arg Gly Ser
 450 455 460
 Val Pro Ile Val Asn Leu Lys Asp Glu Val Leu Ser Pro Ser Trp Asp
 465 470 475 480
 Ser Leu Phe Ser Gly Ser Gln Gly Gln Val Gln Pro Gly Ala Arg Ile
 485 490 495
 Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His Pro Ala Trp Pro Gln
 500 505 510
 Lys Ser Val Trp His Gly Ser Asp Pro Ser Gly Arg Arg Leu Met Glu
 515 520 525
 Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr Gly Ala Thr Gly Gln
 530 535 540
 Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu Gln Lys Ala Ala Ser
 545 550 555 560
 Cys His Asn Ser Tyr Ile Val Leu Cys Ile Glu Asn Ser Phe Met Thr
 565 570 575
 Ser Phe Ser Lys
 580

<210> 17
 <211> 549
 <212> DNA
 <213> Homo sapiens

<400> 17
 cacagccacc ggcacttcca gccggtgctc cacctggttg cgctcaacag cccctgtca 60
 ggcggcatgc ggggcatccg cggggccgac ttccagtgt tccagcaggc gcgggccgtg 120
 gggctggcgg gcaccttccg cgccttccctg tctctcgcc tgcaggacct gtacagcatc 180
 gtgcgccgtg ccgaccgcgc agccgtgccc atcgtcaacc tcaaggacga gctgctgttt 240
 cccagctggg aggtctctgtt ctcaggctct gaggggtccg tgaagcccg ggcacgcac 300
 ttctcctttg acggcaagga cgtcctgagg caccaccct ggccccagaa gagcgtgtgg 360
 catggctcgg accccaacgg gcgcaggctg accgagagct actgtgagac gtggcggacg 420
 gaggtcctc cgccacggg ccaggcctcc tcgctgctgg ggggcaggct cctggggcag 480
 agtgccgcga gctgccatca cgctacatc gtgctctgca ttgagaacag cttcatgact 540
 gcctccaag 549

<210> 18
 <211> 183
 <212> PRT
 <213> Homo sapiens

<400> 18
 His Ser His Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn
 1 5 10 15
 Ser Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln
 20 25 30
 Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ala Gly Thr Phe Arg Ala
 35 40 45
 Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala
 50 55 60
 Asp Arg Ala Ala Val Pro Ile Val Asn Leu Lys Asp Glu Leu Leu Phe
 65 70 75 80
 Pro Ser Trp Glu Ala Leu Phe Ser Gly Ser Glu Gly Pro Leu Lys Pro
 85 90 95
 Gly Ala Arg Ile Phe Ser Phe Asp Gly Lys Asp Val Leu Arg His Pro

100 105 110
 Thr Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Asn Gly Arg
 115 120 125
 Arg Leu Thr Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Ala Pro Ser
 130 135 140
 Ala Thr Gly Gln Ala Ser Ser Leu Leu Gly Gly Arg Leu Leu Gly Gln
 145 150 155 160
 Ser Ala Ala Ser Cys His His Ala Tyr Ile Val Leu Cys Ile Glu Asn
 165 170 175
 Ser Phe Met Thr Ala Ser Lys
 180

<210> 19
 <211> 5408
 <212> DNA
 <213> Homo sapiens

<400> 19
 gccgcgtcga cgcggcggag gaggcagcat cccgcggcgc tgacggtcct ggggagagca 60
 tggcgccgag gtgcccctgg ccatggccgc ggcgggcgcg cctcctggac gtgctcgcgc 120
 ccctggctcct gctgctcggg gtccgcgcgg cctccgcgga gccagagcgc atcagcgagg 180
 aggtggggct gctgcagctc cttggggacc ccccgcccca gcaggtcacc cagacggatg 240
 accccagcgt cgggctggcc tacgtctttg ggccagatgc caacagtggc caagtggccc 300
 ggtaccactt cccagcctc ttcttccgtg acttctcact gctgttcac atccggccag 360
 ccacagaggg cccaggggtg ctgttcgcca tcacggactc ggcgcaggcc atggctctgc 420
 tgggcgtgaa gctctctggg gtgcaggacg ggcaccagga catctccctg ctctacacag 480
 aacctggtgc aggccagacc cacacagccg ccagcttccg gctccccgcc ttcgtcgccc 540
 agtggacaca cttagccctc agtgtggcag gtggctttgt ggccctctac gtggactgtg 600
 aggagttcca gagaatgccg cttgctcgtt cctcacgggg cctggagctg gagcctggcg 660
 ccgggctctt cgtggctcag gcggggggag cggaccctga caagtccag ggggtgatcg 720
 ctgagctgaa ggtgcccagg gacccccagg tgagcccat gactgcctg gacgaggaag 780
 gcgatgactc agatggggca ttccggagact ctggcagcgg gctcggggac gcccgggagc 840
 ttctcagggg ggagacgggc gcggccctaa aaccagggc ccccgcgcca cccccgtca 900
 ccacgcaacc cttggctgga ggcagcagca cgaagattc cagaagtga gaagtgcagg 960
 agcagaccac ggtggcttcg ttaggagctc agacacttcc tggtcagat tctgtctcca 1020
 cgtgggacgg gagtgtccgg acccctgggg gccgcgtgaa agagggcggc ctgaaggggc 1080
 agaaagggga gccagtggtt ccgggcccac ctggcggggc agggcccccga ggatcccat 1140
 gcctacctgg tcccccggtt ctcccggtgc cagtgaagtc cctgggtcct gcaggcccag 1200
 cgttgcaaac tgtccccgga ccacaaggac cccagggccc tccggggagg gacggcaccc 1260
 ctggaaggga cggcgagccg ggcgaccccg gtgaagacgg aaagccgggc gacaccgggc 1320
 cacaaggctt ccctgggact ccaggggatg taggtcccaa gggagacaag ggagaccctg 1380
 gggttgga gaagggggcc ccaggacccc aagggcctcc agggcccccga ggaccctcct 1440
 tcagacacga caagctgacc ttcatgaca tggagggatc tggttttggg ggcgatctgg 1500
 aggcctgcg ggttcctcga ggcttccctg gacctcccg acccccggt gtcccaggcc 1560
 tgcccgcgga gccagccgc tttgggggtga acagctccga cgtcccagga cccgcccggc 1620
 ttctgtgtgt gcctggggcg gaggggtccc ccgggttcc tggtcctccg ggacccccag 1680
 gccctccggg aagagagggg cccccaggaa ggaactgggca gaaaggcagc ctgggtgaag 1740
 caggcgcccc aggcataag gggagcaagg gagcccccg tccctgtggt gctcgtgggg 1800
 agagcgccct ggcaggagcc cccggacctg ctggaccacc agggcccccct gggccccctg 1860
 ggccccagg accaggactc cccgctggat ttgatgacat ggaaggctcc ggggggcctt 1920
 tctggtcaac agcccgaagc gctgatgggc cacagggacc tcccgccctg ccgggactta 1980
 agggggatcc tggcgtgcct gggctgcccg gggcgaaggg agaagtggga gcagatggaa 2040
 tccccgggtt ccccgccctc cctggcagag agggcattgc tgggccccag gggccaaagg 2100
 gagacagagg cagccgggga gaaaaggag atccagggaa ggacggagtc gggcagccgg 2160
 gcctccctgg ccccccgga ccccggggac ctgtgtgcta cgtgtcggag caggacggat 2220

ccgtcctgag	cgtgccggga	cctgagggcc	ggccgggttt	cgcaggcttt	cccggacctg	2280
caggacccaa	gggcaacctg	ggctctaagg	gcgaacgagg	ctccccggga	cccaagggtg	2340
agaagggtga	accgggcagc	atcttcagcc	ccgacggcgg	tgccctgggc	cctgcccaga	2400
aaggagccaa	gggagagccg	ggcttccgag	gacccccggg	tccatacggg	cggccggggg	2460
acaagggaga	gattggcttt	cctggacggc	cgggtcggcc	cgggatgaac	ggattgaaag	2520
gagagaaaag	ggagccggga	gatgccagcc	ttggatttgg	catgaggggg	atgcccgccc	2580
ccccaggacc	tccaggcccc	ccaggccctc	cagggactcc	tgtttacgac	agcaatgtgt	2640
ttgtctgagtc	cagccgcccc	gggcctccag	gattgccagg	gaatcagggc	cctccaggac	2700
ccaaggggcg	caaaggagaa	gtggggcccc	ccggaccacc	agggcagttt	ccgtttgact	2760
ttcttcagtt	ggaggtgaa	atgaaggggg	agaagggaga	ccgaggtgat	gcaggacaga	2820
aaggcgaaaag	gggggagccc	ggggggcgcg	gtttcttcgg	ctccagcctg	cccggccccc	2880
ccggcccccc	aggccacgt	ggctaccctg	ggattccagg	tccaaggga	gagagcatcc	2940
ggggccagcc	cggccacct	ggacctcagg	gacccccggg	catcggtac	gagggggccc	3000
agggccctcc	cggcccccga	ggccccccag	ggcccccttc	atttcctggc	cctcacaggc	3060
agactatcag	cgttcccggc	cctccggggc	ccccggggcc	ccctggggcc	cctggaacca	3120
tggggcgctc	ctcagggggtg	aggctctggg	ctacacgcca	ggccatgctg	ggccagggtc	3180
acgaggttcc	caggggctgg	ctcatcttcg	tggccgagca	ggaggagctc	tacgtccgcg	3240
tgcagaacgg	gttccgggaag	gtccagctgg	aggccccggc	accactccca	caggggacgg	3300
acaatgaagt	ggccgccttg	cagccccccg	tggtgcagct	gcacgacagc	aacccctacc	3360
cgcggcgggg	gcacccccac	cccaccgcgc	ggccctggcg	ggcagatgac	atcctggcca	3420
gccccctcgc	cctgcccag	cccagccct	accccgagc	cccgcaccac	agctcctacg	3480
tgcacctggc	cgcggcgcg	cccacaagcc	cacccgcccc	cagccaccgc	gacttccagc	3540
cgggtgtcca	cctgggttgcg	ctcaacagcc	ccctgtcagg	cggcatgcgg	ggcatccgcg	3600
ggggccgactt	ccagtgtctc	cagcaggcgc	gggcccgtggg	gctggcgggc	accttccgcg	3660
ccttctgtgc	ctcgcgcctg	caggacctgt	acagcatcgt	gcgcggtgcc	gaccgcgcag	3720
ccgtgcccct	cgtcaacctc	aaggacgagc	tgtgttttcc	cagctgggag	gctctgttct	3780
caggctctga	gggtccgctg	aagccccggg	cacgcatctt	ctcctttgac	ggcaaggacg	3840
tcctgaggca	ccccacctgg	cccagaaga	gcgtgtggca	tggtccggac	cccaacgggc	3900
gcaggctgac	cagagcttac	tgtgagacgt	ggcggacgga	ggctccctcg	gccacggggc	3960
aggcctcctc	gctgctgggg	ggcaggctcc	tggggcagag	tgcccgagc	tgccatcacg	4020
cctacatcgt	gctctgcatt	gagaacagct	tcagtactgc	ctccaagtag	ccaccgcctg	4080
gatgcgggat	gcccggagag	accggcggtc	cggagggaagc	ccccaccgtg	ggcagggagc	4140
ggccggccag	ccccggccc	caggacctgg	ctgccatact	ttcctgtata	gttcacgttt	4200
catgtaatcc	tcaagaaata	aaagggaagcc	aaagagtgtg	tttttttaaa	agtttaaaac	4260
agaagcctga	tgtgacatt	cacctgcccc	aactctcccc	tgacctgtga	gcccagctgg	4320
gtcaggcagg	gtgcagtatc	atgccctgtg	caacctcttg	gcctgatcag	accacggctc	4380
gattttctcca	ggatttctctg	ctttgggaag	ccgtgctcgc	cccagcaggt	gctgacttca	4440
tctccacct	agcagcaccg	ttctgtgcac	aaaaccagga	cctgttagca	gacaggcccc	4500
gtgaggcaat	gggagctgag	gccacactca	gcacaaggcc	atctgggctc	ctccagggtg	4560
tgtgtctcgcc	ctgcggtaga	tgggagggag	gctcagggtc	ctggggctag	ggggagcccc	4620
ttctgtctcag	ctctgggcca	ttctccacag	caacccagg	ctgaagcagg	ttcccaagct	4680
cagaggcgca	cgttgacccc	cagctccggc	ctgtcctcca	acaccaagca	cagcagcctg	4740
gggctggcct	cccaaatgag	ccatgagatg	atacatccaa	agcagacagc	tccaccctgg	4800
ccgagtccaa	gctgggagat	tcaagggaacc	catgagttgg	ggctctggcag	cctcccatcc	4860
agggccccca	tctcatgccc	ctggctggga	cgtggctcag	ccagcacttg	tccagctgag	4920
cgccaggatg	gaacacggcc	acatcaaga	ggctgaggct	ggcacaggac	atgcggtagc	4980
cagcacacag	ggcagtgagg	gagggtgtgc	atctgtgcac	tgcccatgga	caggctgggt	5040
ccagatgcag	ggcagtcatt	ggctgtctcc	taggaaaccc	atataccttac	cctccttggg	5100
actgaagggg	aaccccgggg	tgccacagg	ccgcctgcg	ggtgaacaaa	gcagccacga	5160
ggtgcaacaa	ggtcctctgt	cagtcacagc	cacccctgag	atccggcaac	atcaaccgga	5220
gtcattcgtt	ctgtggaggg	acaagtggac	tcagggcagc	gccaggctga	ccacagcaca	5280
gccaacacgc	acctgcctca	ggactgcgac	gaaaccgggt	gggctgggtc	tgtaatgtgt	5340
tgtgatgtga	agccaattca	gacaggcaaa	taaaagtgc	cttttacct	gaaaaaaaaa	5400
aaaaaaaa						5408

<210> 20
 <211> 1336
 <212> PRT
 <213> Homo sapiens

<400> 20

```

Met Ala Pro Arg Cys Pro Trp Pro Trp Pro Arg Arg Arg Arg Leu Leu
 1          5          10          15
Asp Val Leu Ala Pro Leu Val Leu Leu Gly Val Arg Ala Ala Ser
 20          25          30
Ala Glu Pro Glu Arg Ile Ser Glu Glu Val Gly Leu Leu Gln Leu Leu
 35          40          45
Gly Asp Pro Pro Pro Gln Gln Val Thr Gln Thr Asp Asp Pro Asp Val
 50          55          60
Gly Leu Ala Tyr Val Phe Gly Pro Asp Ala Asn Ser Gly Gln Val Ala
 65          70          75          80
Arg Tyr His Phe Pro Ser Leu Phe Phe Arg Asp Phe Ser Leu Leu Phe
 85          90          95
His Ile Arg Pro Ala Thr Glu Gly Pro Gly Val Leu Phe Ala Ile Thr
100          105          110
Asp Ser Ala Gln Ala Met Val Leu Leu Gly Val Lys Leu Ser Gly Val
115          120          125
Gln Asp Gly His Gln Asp Ile Ser Leu Leu Tyr Thr Glu Pro Gly Ala
130          135          140
Gly Gln Thr His Thr Ala Ala Ser Phe Arg Leu Pro Ala Phe Val Gly
145          150          155          160
Gln Trp Thr His Leu Ala Leu Ser Val Ala Gly Gly Phe Val Ala Leu
165          170          175
Tyr Val Asp Cys Glu Glu Phe Gln Arg Met Pro Leu Ala Arg Ser Ser
180          185          190
Arg Gly Leu Glu Leu Glu Pro Gly Ala Gly Leu Phe Val Ala Gln Ala
195          200          205
Gly Gly Ala Asp Pro Asp Lys Phe Gln Gly Val Ile Ala Glu Leu Lys
210          215          220
Val Arg Arg Asp Pro Gln Val Ser Pro Met His Cys Leu Asp Glu Glu
225          230          235          240
Gly Asp Asp Ser Asp Gly Ala Phe Gly Asp Ser Gly Ser Gly Leu Gly
245          250          255
Asp Ala Arg Glu Leu Leu Arg Glu Glu Thr Gly Ala Ala Leu Lys Pro
260          265          270
Arg Leu Pro Ala Pro Pro Pro Val Thr Thr Pro Pro Leu Ala Gly Gly
275          280          285
Ser Ser Thr Glu Asp Ser Arg Ser Glu Glu Val Glu Glu Gln Thr Thr
290          295          300
Val Ala Ser Leu Gly Ala Gln Thr Leu Pro Gly Ser Asp Ser Val Ser
305          310          315          320
Thr Trp Asp Gly Ser Val Arg Thr Pro Gly Gly Arg Val Lys Glu Gly
325          330          335
Gly Leu Lys Gly Gln Lys Gly Glu Pro Gly Val Pro Gly Pro Pro Gly
340          345          350
Arg Ala Gly Pro Pro Gly Ser Pro Cys Leu Pro Gly Pro Pro Gly Leu
355          360          365
Pro Cys Pro Val Ser Pro Leu Gly Pro Ala Gly Pro Ala Leu Gln Thr
370          375          380

```

Val Pro Gly Pro Gln Gly Pro Pro Gly Pro Pro Gly Arg Asp Gly Thr
 385 390 395 400
 Pro Gly Arg Asp Gly Glu Pro Gly Asp Pro Gly Glu Asp Gly Lys Pro
 405 410 415
 Gly Asp Thr Gly Pro Gln Gly Phe Pro Gly Thr Pro Gly Asp Val Gly
 420 425 430
 Pro Lys Gly Asp Lys Gly Asp Pro Gly Val Gly Glu Arg Gly Pro Pro
 435 440 445
 Gly Pro Gln Gly Pro Pro Gly Pro Pro Gly Pro Ser Phe Arg His Asp
 450 455 460
 Lys Leu Thr Phe Ile Asp Met Glu Gly Ser Gly Phe Gly Gly Asp Leu
 465 470 475 480
 Glu Ala Leu Arg Gly Pro Arg Gly Phe Pro Gly Pro Pro Gly Pro Pro
 485 490 495
 Gly Val Pro Gly Leu Pro Gly Glu Pro Gly Arg Phe Gly Val Asn Ser
 500 505 510
 Ser Asp Val Pro Gly Pro Ala Gly Leu Pro Gly Val Pro Gly Arg Glu
 515 520 525
 Gly Pro Pro Gly Phe Pro Gly Leu Pro Gly Pro Pro Gly Pro Pro Gly
 530 535 540
 Arg Glu Gly Pro Pro Gly Arg Thr Gly Gln Lys Gly Ser Leu Gly Glu
 545 550 555 560
 Ala Gly Ala Pro Gly His Lys Gly Ser Lys Gly Ala Pro Gly Pro Ala
 565 570 575
 Gly Ala Arg Gly Glu Ser Gly Leu Ala Gly Ala Pro Gly Pro Ala Gly
 580 585 590
 Pro Pro Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Pro Gly Leu Pro
 595 600 605
 Ala Gly Phe Asp Asp Met Glu Gly Ser Gly Gly Pro Phe Trp Ser Thr
 610 615 620
 Ala Arg Ser Ala Asp Gly Pro Gln Gly Pro Pro Gly Leu Pro Gly Leu
 625 630 635 640
 Lys Gly Asp Pro Gly Val Pro Gly Leu Pro Gly Ala Lys Gly Glu Val
 645 650 655
 Gly Ala Asp Gly Ile Pro Gly Phe Pro Gly Leu Pro Gly Arg Glu Gly
 660 665 670
 Ile Ala Gly Pro Gln Gly Pro Lys Gly Asp Arg Gly Ser Arg Gly Glu
 675 680 685
 Lys Gly Asp Pro Gly Lys Asp Gly Val Gly Gln Pro Gly Leu Pro Gly
 690 695 700
 Pro Pro Gly Pro Pro Gly Pro Val Val Tyr Val Ser Glu Gln Asp Gly
 705 710 715 720
 Ser Val Leu Ser Val Pro Gly Pro Glu Gly Arg Pro Gly Phe Ala Gly
 725 730 735
 Phe Pro Gly Pro Ala Gly Pro Lys Gly Asn Leu Gly Ser Lys Gly Glu
 740 745 750
 Arg Gly Ser Pro Gly Pro Lys Gly Glu Lys Gly Glu Pro Gly Ser Ile
 755 760 765
 Phe Ser Pro Asp Gly Gly Ala Leu Gly Pro Ala Gln Lys Gly Ala Lys
 770 775 780
 Gly Glu Pro Gly Phe Arg Gly Pro Pro Gly Pro Tyr Gly Arg Pro Gly
 785 790 795 800
 Tyr Lys Gly Glu Ile Gly Phe Pro Gly Arg Pro Gly Arg Pro Gly Met
 805 810 815
 Asn Gly Leu Lys Gly Glu Lys Gly Glu Pro Gly Asp Ala Ser Leu Gly

```

      820      825      830
Phe Gly Met Arg Gly Met Pro Gly Pro Pro Gly Pro Pro Gly Pro Pro
      835      840      845
Gly Pro Pro Gly Thr Pro Val Tyr Asp Ser Asn Val Phe Ala Glu Ser
      850      855      860
Ser Arg Pro Gly Pro Pro Gly Leu Pro Gly Asn Gln Gly Pro Pro Gly
      865      870      875      880
Pro Lys Gly Ala Lys Gly Glu Val Gly Pro Pro Gly Pro Pro Gly Gln
      885      890      895
Phe Pro Phe Asp Phe Leu Gln Leu Glu Ala Glu Met Lys Gly Glu Lys
      900      905      910
Gly Asp Arg Gly Asp Ala Gly Gln Lys Gly Glu Arg Gly Glu Pro Gly
      915      920      925
Gly Gly Gly Phe Phe Gly Ser Ser Leu Pro Gly Pro Pro Gly Pro Pro
      930      935      940
Gly Pro Arg Gly Tyr Pro Gly Ile Pro Gly Pro Lys Gly Glu Ser Ile
      945      950      955      960
Arg Gly Gln Pro Gly Pro Pro Gly Pro Gln Gly Pro Pro Gly Ile Gly
      965      970      975
Tyr Glu Gly Arg Gln Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Pro
      980      985      990
Pro Ser Phe Pro Gly Pro His Arg Gln Thr Ile Ser Val Pro Gly Pro
      995      1000      1005
Pro Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Thr Met Gly Ala Ser
      1010      1015      1020
Ser Gly Val Arg Leu Trp Ala Thr Arg Gln Ala Met Leu Gly Gln Val
      1025      1030      1035      104
His Glu Val Pro Glu Gly Trp Leu Ile Phe Val Ala Glu Gln Glu Glu
      1045      1050      1055
Leu Tyr Val Arg Val Gln Asn Gly Phe Arg Lys Val Gln Leu Glu Ala
      1060      1065      1070
Arg Thr Pro Leu Pro Arg Gly Thr Asp Asn Glu Val Ala Ala Leu Gln
      1075      1080      1085
Pro Pro Val Val Gln Leu His Asp Ser Asn Pro Tyr Pro Arg Arg Glu
      1090      1095      1100
His Pro His Pro Thr Ala Arg Pro Trp Arg Ala Asp Asp Ile Leu Ala
      1105      1110      1115      112
Ser Pro Pro Arg Leu Pro Glu Pro Gln Pro Tyr Pro Gly Ala Pro His
      1125      1130      1135
His Ser Ser Tyr Val His Leu Arg Pro Ala Arg Pro Thr Ser Pro Pro
      1140      1145      1150
Ala His Ser His Arg Asp Phe Gln Pro Val Leu His Leu Val Ala Leu
      1155      1160      1165
Asn Ser Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe
      1170      1175      1180
Gln Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ala Gly Thr Phe Arg
      1185      1190      1195      120
Ala Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg
      1205      1210      1215
Ala Asp Arg Ala Ala Val Pro Ile Val Asn Leu Lys Asp Glu Leu Leu
      1220      1225      1230
Phe Pro Ser Trp Glu Ala Leu Phe Ser Gly Ser Glu Gly Pro Leu Lys
      1235      1240      1245
Pro Gly Ala Arg Ile Phe Ser Phe Asp Gly Lys Asp Val Leu Arg His
      1250      1255      1260

```

Pro Thr Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Asn Gly
1265 1270 1275 128
Arg Arg Leu Thr Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Ala Pro
1285 1290 1295
Ser Ala Thr Gly Gln Ala Ser Ser Leu Leu Gly Gly Arg Leu Leu Gly
1300 1305 1310
Gln Ser Ala Ala Ser Cys His His Ala Tyr Ile Val Leu Cys Ile Glu
1315 1320 1325
Asn Ser Phe Met Thr Ala Ser Lys
1330 1335

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/24950

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : A01N 63/00, 43/04; C12N 15/00; C07H 21/02 US CL : 424/93.1; 435/320.1; 514/44; 536/23.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/93.1; 435/320.1; 514/44; 536/23.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	US 5,792,845 A (O'REILLY et al.) 11 August 1998 (11.08.98), col. 4, lines 32-68, col. 5, lines 1-2, 51-68, col. 6, lines 1-8.	1-30, 33
Y	WO 97/23500 A1 (THE CHILDREN'S MEDICAL CENTER CORPORATION) 03 July 1997 (03.07.97), page 41, lines 3-33, page 42, lines 1-27.	4
X,P	WO 98/49321 A2 (RHONE-POULENC RORER) 05 November 1998 (05.11.98), page 44, 6-11, 25-33, page 45, lines 12-13, 29-35.	1, 5, 18, 20, 31
Y,P		2-4, 6-17, 19, 21-30, 32, 33
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:	* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*a* document member of the same patent family	
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 04 FEBRUARY 1999	Date of mailing of the international search report 08 MAR 1999	
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer SHIN-LIN CHEN	
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196	

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US98/24950

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97/15666 A (THE CHILDREN'S MEDICAL CENTER CORPORATION) 01 May 1997 (01.05.97), page 20, lines 16-35, page 21, page 22, lines 1-16. page 59, lines 5-35, page 60, page 61, 1-6.	1-30, 33
Y	TANAKA, T. et al. Retroviral and adenoviral mediated transduction of angiostatin cDNA inhibits angiogenesis and tumor growth. Proceedings of the American Association for Cancer Research. March 1997 (03.97). Vol 38. page 264.	1-33
Y	WO 96/35774 A2 (THE CHILDREN'S MEDICAL CENTER CORPORATION) 14 November 1996 (14.11.96), page lines 33-36, pages 22-25, page 26, lines 1-33. pages 144-148.	1-30, 33
Y	WO 97/41824 A2(ABBOTT LABORATORIES) 13 November 1997 (13.11.97), page 5, lines 13-38, page 6, 1-18, page 60, lines 15-38, pages 61-62, page 63, lines 1-33.	1-30, 33
Y	WO 95/29242 A1 (THE CHILDREN'S MEDICAL CENTER CORPORATION) 02 November 1995 (02.11.95), page 21, lines 19-35, pages 22-27, page 28, lines 1-4. page 87, lines 4-35, page 88, page 89, lines 1-14.	1-30, 33
Y	O'REILLY, et al. Angiostatin induces and sustains dormancy of human primary tumors in mice. Nature Medicine, June 1996 (06.96), Vol. 2, No. 6, pages 689-692, especially pages 689-690.	1-30, 33
Y	O'REILLY, et al. Endostatin: An Endogenous Inhibitor of Angiogenesis and Tumor Growth. Cell, 24 January 1997 (24.01.97), Vol. 88, pages 277-285, especially pages 279-280, 282.	1-30, 33

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/24950

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, STN, WPIDS, MEDLINE, CAPLUS, BIOSIS, EMBASE

search terms: angiostatin, plasminogen, endostatin, collagen(w) XVIII, inhibit?(5a)tumor(5a)growth, tumor(5a)regress?,
diabet?(p)retinopathy, plasmid, viral(5a) vector.